



COMPREHENSIVE COMMUNICABLE DISEASE MANAGEMENT PLAN

Adapted for MESD and its component districts (Reynolds School District) with permission from Dr. Jan Olson and team's comprehensive communicable disease management plan for the Molalla River School District

THIS PLAN CONTAINS:

RSD Communicable Disease Prevention Plan

RSD Exposure Control Plan

RSD Pandemic Response Plan

COVID-19 Specific Considerations

TABLE OF CONTENTS

INTRODUCTION	3
COMMUNICABLE DISEASE PREVENTION PLAN	4
Communicable Disease Prevention	5
Common Childhood Infectious Diseases	5
Vaccines	5
Hygiene	6
Respiratory Hygiene/Cough Etiquette	7
Environmental Surface Cleaning	7
Communicable Disease Exclusion	7
Restrictable Diseases	8
Isolation Spaces	8
Outbreaks of Illness, and Symptom Clusters	9
Respiratory Illness	9
Vaccine Preventable Disease	9
Gastroenteritis	10
Other Circumstances	10
Animals in School	10
Food Safety	11
EXPOSURE CONTROL PLAN	12
Exposure Prevention	13
Employee Exposure Determination	13
Universal & Standard Precautions	14
Hand Hygiene	14
Personal Protective Equipment (PPE)	14
Respiratory Hygiene/Cough Etiquette	15
Engineering and Work Practice Controls	16
Regulated Biohazardous Waste Containers	16
Sharps Safety	16
Health Room, Isolation Space, and Classroom Safety	17
Cleaning and Disinfecting Environmental Surfaces	17
Transmission-Based Precautions	18
Contact Precautions	18
Droplet Precautions	18
Airborne Precautions	18
Body Fluid Exposure Incidents	19
Needle-Stick Injuries	19
Mucous Membranes	19
Bites	19
MESD Body Fluid Exposure Protocol	20
PANDEMIC PLAN	21
Control Measures	21
Special Considerations	25
GLOSSARY & REFERENCES	26
COVID-19 SPECIFIC COMMUNICABLE DISEASE MANAGEMENT ADDENDUM	29
COVID-19 TOOLS	44

INTRODUCTION

The health and safety of all students and staff is a priority of Multnomah Education Service District (MESD). The control of communicable diseases is an essential component of health and wellness in the school setting. Providing a safe, comfortable, and healthy environment facilitates the educational process, encourages social development, and allows children to develop healthy attitudes toward school (NRC, 2020).

Illness and injury are not uncommon in the school setting and thus policies, procedures, and guidance regarding infection control are of utmost importance. When children are injured or feel unwell at school, it can create risk to others and impact the ability of a child to fully participate in their educational activities. In the Whole School, Whole Community, Whole Child model, school personnel collaborate for the best outcomes of the individual student, and the school population as a whole. To accomplish this goal, staff must have access to and be trained on resources and materials to identify appropriate measures and interventions for child health issues (ACSD, 2020).

The purpose of this comprehensive guide is to provide infection control guidance, practice standards, and protocols for Multnomah Education Service District (MESD) and its component school districts (Reynolds School District).

This document combines the district's *Communicable Disease Prevention Plan*, *Exposure Control Plan*, and *Pandemic Response Plan* to form a Comprehensive Communicable Disease Management Plan.

This plan was adapted (with permission) from Dr. Jan Olson and team's comprehensive communicable disease management plan for the Molalla River School District, by the MESD School Health Services department in collaboration with district administration.

Original Author: Dr. Jan Olson, DNP, MSNEd, BSN, RN, NCSN, District Nurse

Original Collaborators: Amanda Bickford, BSN, RN, District Nurse

Tony Mann, Superintendent Rick Gill, Risk Manager

Dave Luce, Human Resources Administrator

Original Date: June 3, 2020

MESD Adaptation Adoption Date: July 13, 2020

Adaptation Collaborators: Dr. Paul Coakley, MESD Superintendent

Jamie Smith, MPH, BSN, RN, NCSN, Coordinator of School Health Services – MESD

Lisa Ferguson, MS, BSN, RN, Communicable Disease Manager - Investigations and Case Management - Multnomah County Health Department

Heather Godsey, BSN, RN, Nurse Consultant – MESD School Health Services

Angela Kulak, BSN, RN, Nursing Supervisor – MESD School Health Services

Lorraine Parga, BSN, RN, Nursing Supervisor – MESD School Health Services

Chandra Siim, BSN, RN, Nurse Consultant – MESD School Health Services




**This manual, or portions of it, may not be reprinted without permission of School Health Services, Multnomah Education Service District. The EXPOSURE CONTROL PLAN in this manual is published by MESD to provide information on the development of an appropriate Exposure Control Plan. The manual is not intended as a substitute for a district's own careful review of OR-OSHA regulations which should be the basis for plan development. Agencies/Districts contracting with School Health Services for Bloodborne Pathogens Training may copy and/or revise all forms within this manual to meet their individual needs.*

COMMUNICABLE DISEASE PREVENTION PLAN

Communicable disease control and prevention is of significant importance in creating a safe and healthy environment for students and staff.

A communicable disease is an infectious illness that is transmissible by contact with:

- Bodily discharge or fluids of an infected individual
Contact with contaminated surfaces or objects
- Contact with contaminated surfaces or objects
- Inhalation of respiratory droplets expelled by an infected individual
- Ingestion of contaminated food or water
- Direct or indirect contact with disease vectors



Although the terms *communicable disease*, *contagious disease*, and *contagious condition* are often used interchangeably, it is important to note that not all communicable diseases that are spread by contact with disease vectors are considered to be “contagious” diseases since they cannot be spread from direct contact with another person (ACPHD, 2013). In the school setting there is a prevention-oriented approach for communicable disease management which is grounded in education, role modeling, and implementation of standard precautions and hygiene. However, the nature of a population-based setting lends to the need to establish practices for measures and interventions associated with exposures or potential exposures. This first section focuses on a population-based set of practices for communicable disease prevention, while the subsequent Exposure Control Plan discusses work-practice control measures for staff, per the [OSHA Bloodborne Pathogen Standard](#) (BBP, 29 CFR 1910.1030).

RSD Board Policies

[Communicable Diseases JHCC](#)

[Communicable Diseases – Staff GBEB-AR](#)

[Communicable Diseases – Staff GBEB](#)

[Communicable Diseases – Students JHCC](#)

[Student Health Services and Requirements JHC](#)

MESD Board Policies

[Communicable Diseases JHCC/EBBA/GBEB-AR](#)

[School Health Services and Requirements JHC](#)

Oregon Legislation

[OAR 333-019-0010 Disease Related School, Child Care, and Worksite Restrictions: Imposition of Restrictions](#)

[OAR 581-022-2200 Health Services](#)

[ORS 410-133-0000 School Based Health Services](#)

Oregon Health Authority & Oregon Department of Education

[Oregon Department of Education Communicable Disease Guidelines for Schools](#)

Communicable Disease Prevention

There are many methods that can be applied to control communicable diseases at a variety of levels. Some of the most common include vector control, hygiene, sanitation, and immunization. Fully endorsing the control and prevention of communicable diseases requires an understanding of disease transmission.



Hand Hygiene



Cough Etiquette



Immunizations



Blood Borne Pathogen Training



Environmental Sanitation



Standard Precautions



Illness Policy



Food Safety

How these communicable diseases are spread depends on the specific infectious agent. Common ways in which communicable diseases spread include:

- Physical contact with an infected person, such as through touch (staphylococcus), sexual intercourse (gonorrhea, HIV), fecal/oral transmission (hepatitis A, C. difficile), or respiratory droplets (influenza, varicella-zoster)
- Contact with a contaminated surface or object (norovirus), food (salmonella, E. coli), blood (HIV, hepatitis B, hepatitis C), or water (cholera, listeria)
- Bites from insects or animals capable of transmitting the disease (mosquito: malaria and yellow fever; flea: plague)
- Inhalation of small respiratory droplets that are dispersed through or suspended in the air (measles, tuberculosis)

In the school setting the most frequent risks are associated with direct contact with ill individuals, contaminated surfaces, or inhalation of infectious droplets. Primary methods of illness prevention include hand hygiene, sanitation respiratory etiquette, isolation and exclusion of symptomatic individuals, as well as standard and transmission-based precautions.

This section of the plan will provide a brief overview of, and procedures for addressing the following communicable disease issues in the school setting:

- Common Childhood Infectious Diseases
- Vaccines
- Hand Hygiene
- Respiratory/Cough Etiquette
- Environmental Surface Cleaning

[The district Exposure Control Plan in this manual discusses Standard Precautions in detail as well as Transmission Based Precautions which include contact, droplet, and airborne precautions. The District Pandemic Plan will address measures specific to novel virus response.]

Common Childhood Infectious Diseases

There are a variety of [Common Childhood Infectious Diseases](#) that are regularly encountered in the school setting. Some childhood illnesses such as the common cold (adenoviruses, coronaviruses, rhinoviruses), bronchitis, sinusitis, and tonsillitis occur throughout the year. While other illnesses such as gastroenteritis (norovirus most frequently), croup (most commonly parainfluenza), and influenza (A & B) most often occur seasonally. Examples of other common childhood infections include: strep throat, hand foot and mouth disease, fifth disease, and staph skin infections. There are other more severe infectious diseases, such as bacterial meningitis, that also occur sporadically throughout the school year (BCDC, 2009).

Vaccines

In the school setting, vaccines are an important step toward communicable disease control and certain vaccines are required for attending school in Oregon. However, it is important to note that some individuals may not be vaccinated due to medical/non-medical exemptions. Each school has a record of which students are and are not vaccinated with routine childhood immunizations as a primary control measure for outbreaks of vaccine preventable diseases. You can find [more information on immunizations](#) on the [MESD School Health Services Website](#).

Under direction of the MESD Communicable Disease Team:

- When a positive case of a vaccine preventable disease (varicella, pertussis) is identified in the school setting, designated staff will run immunization reports to identify unvaccinated students in the building. The Communicable Disease Team will work with the local public health authority and the school administrator to determine the need for exclusion or notification of exposure to members of the school community.
- In the event that a positive case is identified in the school building, or when the circulation of a vaccine preventable disease (such as measles) is increasing in incidence in a building or in the community, the nurse consultant will work with the local public health authority to determine the necessity for exclusion of unvaccinated students or staff.


Hygiene

Prevention-oriented measures are grounded in disease transmission education, and the practical application of general sanitation and personal hygiene. Hygiene and sanitation are some of the most important methods of disease prevention. Handwashing is the single most important method of keeping germs at bay, specifically in the school setting. Appropriate handwashing practices should be taught, modeled by staff, and practiced by all.

[Age appropriate hand hygiene curriculum](#) can be found from a variety of resources and should be provided annually in the fall and as needed during peak illness season or specific increases of disease in the school setting.

Hand sanitizer, while not effective against a large number of pathogens, should be made available for times that handwashing is not immediately accessible. Hand sanitizer, that contains at least 60% alcohol, should be easily accessible throughout the building, specifically in high contact areas, such as entrances/exits, water fountains, cafeterias, gyms and classroom. Supervised stocks of hand sanitizer should be available in each classroom.

How to wash your hands



Students and staff should wash hands:

- Before, during, and after preparing food
- Before eating food
- Before and after caring for someone at home who is sick with vomiting or diarrhea
- Before and after treating a cut or wound
- After using the toilet
- After changing diapers or cleaning up a child who has used the toilet
- After blowing your nose, coughing, or sneezing
- After touching an animal, animal feed, or animal waste
- After handling pet food or pet treats
- After touching garbage
- Before and after using a hand-operated water fountain
- Before and after using PPE (gloves, mask)
- When hands are visibly soiled

(CDC, RSSL, 2022)

When immunocompromised students and staff are present, increase in hand hygiene frequency is a necessary illness-prevention intervention.

Respiratory Hygiene/Cough Etiquette

Respiratory hygiene and cough etiquette are terms used to describe infection prevention measures to decrease the transmission of respiratory illness (e.g., influenza and cold viruses). A respiratory infection can be spread when a person who is infected coughs or sneezes. The droplets released from an ill person's cough or sneeze can travel for several feet, reaching the nose or mouth of others, causing illness. Pathogens can spread easily from person to person through direct contact via touching objects or shaking hands, and then touching one's eyes, nose, or mouth. Droplets can persist for a short time on a variety of objects and high-contact areas such as doorknobs or desks.

Some individuals cough without having respiratory infection (e.g., persons with chronic obstructive lung disease), and because some individuals can carry and spread diseases without ever showing any symptoms of illness. As a result, we do not always know who is infectious and who is not. Therefore, respiratory hygiene and cough etiquette are very important components to protecting yourself from illness and preventing others from becoming ill.

Like hand hygiene, respiratory hygiene is part of the standard precautions that should be taught, practiced, and modeled to prevent the spread of disease. (See Respiratory Hygiene and Cough Etiquette for more detailed information on practices and interventions)



(Image: Manitoba Department of Health)

Environmental Surface Cleaning

Clean schools contribute to healthy environments and reduce the risk of communicable disease transmission. Some of the important concepts associated with reduction in illness include:

- Scheduling *and documenting* the routine cleaning of classrooms, common areas, and shared objects and surfaces;
- Ensuring adequate stock of appropriate sanitizers and disinfectants;
- Ensuring garbage and sharps containers are emptied before they become full;
- Ensuring that any classrooms with pets have a cleaning plan in place to minimize odors or contamination.

While environmental cleaning is largely governed by facilities management and custodial services, there are certain classroom measures that can be practiced to improve cleanliness and reduce the risk of illness-transmission during peak illness times, such as increasing access to sanitizing wipes, tissue, and hand sanitizer. (See the *Exposure Control Plan* for more detail on disinfection.)

Communicable Disease Exclusion *(See COVID-19 Addendum for additional exclusion guidance during the COVID-19 pandemic)*

In the school environment, communicable diseases are easily transmitted from one individual to another by various routes, and can even be transmitted while an individual is not showing symptoms of illness. While some conditions are restrictable based on diagnosis, more often early identification of signs and symptoms of communicable disease is of paramount importance to increase the health of the school population and decrease school absenteeism. Effective control measures include education, avoidance of risk factors, sanitation, air filtration/ventilation, vaccination, early recognition of symptoms, health assessment, prompt diagnosis, and adequate isolation and treatment (ODE, 2020).

Oregon public health law (Oregon Administrative Rule 333-019-0010) mandates that persons who work in or attend school who are diagnosed with certain diseases or conditions be excluded from school until no longer contagious. Since diagnosis often presumes assessment by a licensed health care provider and specific testing, schools must often make decisions regarding exclusion based on non-diagnostic but readily identifiable symptoms. As a matter of routine practice, individuals with the following excludable symptoms of illness should be excluded from school per OAR 333-019-0010, Local Public Health Authority (LPHA) guidelines, and the [Oregon Department of Education Communicable Disease Guidance Document](#).

Restrictable Disease

Restrictable diseases are specific infectious disease diagnoses that require students or staff to remain at home for a specified amount of time to limit transmission. Restriction is typically associated with the communicability or severity of a disease. Some restrictable diseases are also *reportable to the LPHA, in which case the MESD Communicable Disease Team should always be contacted. The LPHA typically also notifies the MESD Communicable Disease Team of confirmed cases of reportable illness in a student or staff member. There are occasions when a parent/guardian will notify the school of a communicable disease first. In these cases the school nurse (RN) should be notified and they will consult with the MESD Communicable Disease Team.

Students with diagnoses of diseases restrictable by the LPHA under OAR 333-019-0010 should return to school when documentation is obtained from the LPHA or healthcare provider indicating they are no longer infectious. This includes, but is not limited to:

- COVID-19*
- Diphtheria*
- Hepatitis A*
- Hepatitis E*
- Measles*
- Mumps*
- Pertussis* (whooping cough)
- Rubella*
- Salmonella enterica serotype Typhi infection*
- Scabies
- Shiga-toxigenic Escherichia coli (STEC) infection*
- Shigellosis*
- Varicella zoster (chickenpox)
- Infectious tuberculosis*

- » If a report is made to the school office, administration, or other school staff regarding any communicable disease diagnosis in students or staff, **this should immediately be referred to the school nurse** who will then consult with the MESD Communicable Disease Team.
- » This should be regarded as an urgent referral to the MESD Communicable Disease Team if the disease is regarded as a **reportable** condition.
- » **School staff receiving reports should follow confidentiality rules by not informing any other students, staff, or parent(s)/guardian(s) of the report, besides notifying the administrator and school nurse.**
- » The MESD Communicable Disease Team will confirm the diagnosis with the LPHA, and identify the need for communication, surveillance, or control measures. The interventions and communication are driven by multiple factors including the diagnosis, student health status, risk of exposure, number of individuals infected, and risk to cohorts or specific students and staff.
- » Depending on the diagnosis and the identified interaction with a reported case, the MESD Communicable Disease Team may ask that the school nurse directly notify any individuals who are at high-risk of severe disease, immunocompromised or pregnant, so that they can reach out to their healthcare provider with any questions or concerns regarding a potential exposure. The school nurse will work with the school' administrator to ensure that any staff members or students who have identified as being in this higher-risk category will receive immediate notification of potential exposures.

Isolation Spaces

Oregon Law (OAR 581-022-2220) requires the school district to maintain a prevention oriented health services program which includes health care and space that is appropriately supervised and adequately equipped for first aid, and **isolates the sick or injured child from the student body.**

When students are identified with restrictable diseases or excludable symptoms, they should be separated from the well-population, in an appropriate space until they can be dismissed to home. This isolation space should be separated from the healthcare area used to assess and treat injured and non-symptomatic children or to provide medication management and care of chronic health conditions.

Outbreaks of Illness, and Symptom Clusters

Outbreaks are most often defined as compatible diagnoses or syndromes in individuals from 2 or more households that are epidemiologically-linked and in the same communicability time period. Because of the nature of the ongoing congregating setting of school, this definition is insufficient for the purposes of seasonal illness; rather an increase in morbidity or severity should be indicators to report to the school nurse and the MESD Communicable Disease Team for consideration of outbreak reports or control measure implementation. The attention to outbreaks, interventions, and resources are highly dependent on the prevalence, severity and communicability of the disease. Outbreak investigations will be facilitated through the MESD Communicable Disease Team, in collaboration with the school nurse and health assistant, the school or district administrator, and the LPHA with the use of the [Oregon Health Authority Outbreak Toolkits for Schools](#).

Respiratory Illness

Respiratory illness or disease refers to the pathological conditions affecting the organs and tissues that make gas exchange possible, including conditions of the upper respiratory tract, trachea, bronchi, bronchioles, alveoli, pleura and pleural cavity, and the nerves and muscles of breathing. Respiratory diseases range from mild and self-limiting, such as the common cold, to life-threatening, like bacterial pneumonia.

Individuals with respiratory illnesses are often present in the school setting. The following indicators should be reported to the school nurse, who will consult with the MESD Communicable Disease Team:

- Any respiratory illness resulting in hospitalization or death of a student or staff member
- Diagnosed pneumonia in 3 or more individuals in the same cohort
- Unusually high levels of absenteeism due to respiratory illnesses ($\geq 30\%$ of the student/staff with at least 10 individuals absent) at the school level on any given day.
- Unusually high levels of absenteeism due to respiratory illnesses ($\geq 20\%$ of a cohort with at least 3 individuals absent) within the cohort level on any given day
- Prolonged illness, lasting longer than 3 days on average, among 10 or more persons (or $\geq 20\%$, whichever is greater) in the same cohort

*In the event of respiratory illnesses related to novel viruses, the Pandemic Plan will be deferred to.

Vaccine Preventable Disease

A vaccine-preventable disease (VPD) is an infectious disease for which an effective preventive vaccine exists. Current VPDs routinely immunized for in the United States include:

- Diphtheria*
- Tetanus*
- Measles*
- Mumps*
- Rubella*
- Haemophilus influenzae type b infections (Hib)*
- Pneumococcal infections*
- Meningococcal disease*
- Pertussis (whooping cough) *
- Poliomyelitis (polio)*
- Hepatitis A*
- Hepatitis B*
- Varicella (chickenpox)
- Influenza

Most VPD's are also reportable diseases*, meaning they are under constant surveillance and the LPHA should be notified of any case. Some communicable diseases are uncommon locally, and are not routinely immunized for in the US, so a diagnosed case would be of interest. This may include:

- Cholera
- Plague
- Rabies
- Bat lyssavirus
- Yellow fever
- Japanese encephalitis
- Q fever
- Tuberculosis
- Typhoid

Uncommon diseases may also pose a risk to a susceptible individual or population if circumstances permit exposure and should therefore also be reported to the LPHA.

Reports of VPDs should be referred to the MESD Communicable Disease Team, whether coming from a parent, provider, community member, or the LPHA. Indicators for possible reporting to the LPHA of school-identified VPDs include:

- A single case of a vaccine preventable disease that is also a reportable disease or uncommon locally
- More than 2 cases of diagnosed chickenpox (varicella) from separate households in the same cohort or more than 5 cases in a school.
- More than 3 cases of diagnosed influenza from separate households in the same cohort within 72 hours.

Gastroenteritis

An outbreak of gastroenteritis is defined as more cases than expected for a given population and time period. For example, two children in a 25-person classroom with vomiting or diarrhea within one week could potentially indicate an outbreak. Because the nature of viral gastroenteritis (norovirus) is common, seasonal, and highly infectious, it is unlikely to result in an outbreak investigation unless the number infected and/or duration is unusual. However, preventive measures to reduce spread would need to be immediately enhanced and enforced. Because symptoms of bacterial gastroenteritis may start with a similar presentation, it is important to evaluate the severity for the duration of illness.

Gastrointestinal indicators to report to the school nurse include:

- Multiple children with compatible symptoms in 48 hours within the same cohort, but separate households
- More than 2 cases of diarrhea with bloody stool in the school setting
- Sudden onset of vomiting in multiple persons (from separate households) in the same cohort
- Any unusual combination of gastrointestinal symptoms, severity, duration, or incidence

Other Circumstances

While less common, outbreaks of skin infections, other novel diseases, or unusual infectious disease circumstances can arise.

To ensure the appropriate disease intervention, surveillance and data collection occur, these unusual situations should be referred to the MESD Communicable Disease Team, and will be handled on a case by case basis. Examples of these circumstances may include:

- More than 2 students from separate households with reported compatible skin infections in the same cohort or athletic team
- Any student or staff member coming into contact with blood, saliva, or feces from a non-domestic animal
- Any staff coming into contact with blood or other potentially infectious body fluid that is not their own (following the MESD Exposure Control Plan); *students coming into contact with the body fluids of others should be referred to the school nurse or administrator*
- Any unusual combination of gastrointestinal symptoms, severity, duration, or incidence
- Any combination of illness symptoms, severity, duration, or frequency that seems unusual as compared to routine seasonal illness

In regard to determined outbreaks or novel diagnoses the MESD Communicable Disease Team may decide that additional control measures or data collection are necessary, and will consult with the building administrator and the LPHA as needed.


The MESD Communicable Disease Team should always be consulted regarding any written communication that may be developed to notify parents, students, or staff about illness, disease outbreaks, and risks to students, families, and staff, and/or control measures specific to an outbreak or exposure. Any presentation of illness or combination of illnesses as described above should be reported to the MESD Communicable Disease Team and the building administrator.

Animals in School

Animals in schools can have a positive effect in the school environment, but also may cause infectious disease issues or other issues for staff and students. MESD only allows for animals on district property with specific approval under certain circumstances. School board policies and district applications should be visited for clarification on this. Other considerations should be made in regard to controlling spread of infectious diseases from animals:

- Wild mammals, alive or recently dead, should not be allowed in the school setting. Bats and skunks have a significant risk of being rabid. Other wild animals may be more prone to causing injury through bites and scratches.
- Dogs, cats, and ferrets allowed in school buildings should have a current rabies vaccine.

- Any animal bites on school property should be reported to the LPHA for follow up. Standard first aid should be provided and the individual should be referred to their health care provider.
- Animals who are ill should not be allowed into the school setting.
- Class pets should be removed if they become ill.
- Handwashing must occur before and after handling of animals to prevent disease transmission.
- Animals should not be present or handled in areas where food and drink are consumed or prepared.
- Children should not kiss high risk animals such as chicks, ducks, turtles, and other reptiles.
- Children should always be monitored during animal interactions.
- Consider the medical needs of students who are immunosuppressed or who have allergies, as they may become severely ill when exposed to certain pathogens or allergens.
- In the event that a student in a classroom is diagnosed with a disease known to be carried by animals (i.e., campylobacteriosis or salmonellosis), for example) any classroom animals should be removed from the classroom setting until the risk is determined to be resolved.



Food Safety

Food safety training and enforcement for nutrition services staff is supervised by nutrition services. For the overall wellness of our school based population, all food preparation and consumption within the classroom should follow general food safety standards and disease prevention principles.

For elementary school classrooms

- Hand hygiene should be enforced prior to eating
- General principles of food safety can be taught that are age appropriate
- Food sharing should not be allowed
- For classroom and school sponsored events, only commercially prepared products are permitted. Do not allow homemade food from non-licensed kitchens to be served.

For middle school or high school culinary classrooms

- Hand hygiene should always be enforced
- Age appropriate food safety principles are taught
- Appropriate food handling processes must be taught, role modeled, and enforced. This includes overview of:
 - Hand hygiene and appropriate use of gloves
 - Clean surfaces and appropriate use of sanitizers
 - Separating raw and ready to eat foods/ avoidance of cross contamination
 - Cooking food to appropriate temperatures
 - Appropriate storage and refrigeration
 - Measures to prevent allergic reactions
 - Abstaining from food preparation when specific symptoms or specific illnesses have been identified



EXPOSURE CONTROL PLAN

This plan provides the employees of MESD with guidelines for handling any exposure to blood or other potentially infectious materials (OPIM). These established procedures are in accordance with local and state requirements, as well as federal occupational safety and health requirements (OSHA BBP Standard 1910.1030). Districts that contract with MESD for Bloodborne Pathogens training for their at-risk staff are assumed to agree to and ensure compliance with this Exposure Control Plan unless they specifically notify the MESD of any variance from the plan, in writing.

The principle of Standard Precautions holds that **all body fluids or other potentially infectious materials should be considered potentially infectious at all times, since many disease-causing bacteria and viruses may be carried in the body fluids of persons who show no symptoms of illness**. Standard Precautions shall be observed in all MESD sites in order to prevent contact with all body fluids and other potentially infectious materials, including blood, semen, vaginal secretions, cerebrospinal fluid, synovial fluid, pleural fluid, pericardial fluid, peritoneal fluid, amniotic fluid, saliva in dental procedures, any body fluid that is visibly contaminated with blood, and all body fluids in situations where it is difficult or impossible to differentiate between body fluids. Transmission-based Precautions should additionally be endorsed in circumstances where specific risk is anticipated based on health status.

Disease transmission is facilitated by three conditions: high infectivity of a body fluid from an infectious person (carrier), a portal of exit from the carrier, and a portal of entry into a susceptible person. Any staff who may be expected to clean up body fluid spills, perform personal care for a student, perform initial first aid or injury response, or who are working with populations less able to control their bodily fluids in a congregate setting such as a school, are reasonably anticipated to have “occupational exposure” to blood or other potentially infectious material. These staff should be provided the required training and Personal Protective Equipment (PPE) so that they are aware of their risks and are able to best protect themselves.

RSD Board Policies

[Infection Control – HIV, AIDS, HBV EBBAA/GBEBC/JHCCC](#)

[Bloodborne Pathogens – Exposure Control Plan for Compliance with OR-OSHA Standard GBEBC/JHCCC/EBBAA-AR](#)

[Staff Development – Nonlicensed GDL-AR](#)

[Communicable Diseases JHCC/EBBA/GBEB-AR](#)

[Communicable Diseases – Students and Staff JHCC/GBEB](#)

MESD Board Policies

[Infection Control – HIV, AIDS, HBV EBBAA/GBEBC/JHCCC](#)

[Bloodborne Pathogens – Exposure Control Plan for Compliance with OR-OSHA Standard GBEBC/JHCCC/EBBAA-AR](#)

[Staff Development – Nonlicensed GDL-AR](#)

[Communicable Diseases JHCC/EBBA/GBEB-AR](#)

[Communicable Diseases – Students and Staff JHCC/GBEB](#)

OSHA

[Bloodborne Pathogens 1910.1030](#)

[Personal Protective Equipment 1910 Subpart 1](#)

Exposure Prevention

Per OSHA regulations, in order to reduce risk and promote prevention of infections related to blood or body fluids, the district will provide or promote specific trainings or practices to prepare staff, including:

- Bloodborne Pathogens (BBP) Training, an **annual** requirement for all at-risk staff. **The first class must be in-person**, and the subsequent annual updates may be taken online **through MESD's website** during work hours (as long as the tasks to which the employee is assigned do not change). **Please note that the Safe Schools/Vector BBP Module, while a good general course for all school staff to take, is **not** an OSHA-approved training which meets the requirements for the training of at-risk staff.*
- Hepatitis B vaccination (education and recommendations on Hepatitis B vaccination are provided each year with BBP training). A waiver may be signed in lieu of immunization if staff opt out AFTER completing in-person BBP training and acknowledging their understanding of the risk and implications.
- Consistent use of Standard Precautions is expected any time the risk of exposure to body fluids is present.
- Routine training, refreshers, and understanding of appropriate first aid.
- Routine training or refreshers for staff who provide direct care to students or who work with students with specific disabilities, or health or behavior plans.

Employee Exposure Determination

Any employee with a risk of occupational exposure to blood and other potentially infectious body fluids is protected by the OSHA-required EXPOSURE CONTROL PLAN. Exposure determination for common job classifications in schools are noted below. Exposure determination of school employees is the responsibility of the school or district administrator. School districts must review job tasks and make individual determinations regarding exposure risks based on OR-OSHA criteria. Each school environment will need an individual assessment of risks to various employees. School administrators may need assistance with determination of risk for certain employees due to their specific job tasks and are encouraged to seek consultation with the MESD, the district risk management department, or legal counsel regarding this risk of exposure.

CLASS A EXPOSURES: Job classifications in which employees are assumed to have occupational exposure due to direct physical care in which blood or other potentially infectious material could be present (such as direct care for injuries, illness, medical/dental procedures, injections, etc.):

- School Health Services staff in health-rooms or isolation spaces
- Secretaries or other office assistants in schools without available health services staff
- Custodial staff or any other staff expected to clean up body fluid spills
- Athletic Trainers/Coaches/P.E. Teachers dealing with acute injuries
- Unlicensed Assistive Personnel (UAP) who provide tasks taught or delegated by the school nurse
- Staff working with SpEd populations or other populations who are not able to adequately contain their body fluids, or who expose others through behaviors such as biting, spitting, fecal smearing, and scratching.
- Staff who have a job description which requires them to provide first aid to students/staff

CLASS B EXPOSURES: Job classifications in which employees may be expected to have *some* occupational risk:

- General Educational Teachers/Educational Assistants & Substitutes
- School Psychologist
- Physical/Occupational Therapists
- Speech Language Pathologists
- Students and Instructors in Health Occupation Programs
- Bus Drivers

****It is the responsibility of the individual school district administrator to determine which school employee job descriptions place employees at risk for body fluid exposure.***

Universal & Standard Precautions

The premise of Universal Precautions is to treat all body fluids as potentially infectious. Standard Precautions align with this and provide a set of standards for hygiene and barrier protection (or PPE) during any and all encounters with body fluids.

Standard Precautions are regarded as the minimum infection prevention practices that apply to all direct care or exposure to body fluids, regardless of suspected or confirmed infection status of the individual, in any setting where there is an expected risk of body fluid exposure. In the school setting, body fluid exposures most frequently occur with physical injury. Exposures may also occur relative to a health related issue, medical procedure, developmental issue, behavioral issue or disability. Students and co-workers should be encouraged to care for their own bleeding injuries. If assistance is necessary, the use of disposable gloves and other barriers, followed by hand washing, is required for the designated caregiver when body fluids are present. There may be occasions when unanticipated contact with body fluids will occur, and in such cases, staff should follow the MESD Body Fluid Exposure Protocol (to be discussed below).



Standard Precautions endorse the appropriate use of (PPE and other practices such as hand hygiene and respiratory etiquette, as well as work practice controls such as sharps safety (handling needles, lancets, broken glass, etc.) and environmental disinfection.

When Standard Precautions alone cannot prevent transmission, they are supplemented with Transmission-Based Precautions. This second tier of infection prevention is used when there is a specific risk related to an ill student or staff in the school setting that can spread through contact, droplet, or airborne routes (e.g., skin contact, sneezing, coughing) and are always used in addition to Standard Precautions. While use of Transmission-Based Precautions are typically restricted to the health room/isolation space under specific conditions, there is still a risk of exposure in the school setting.

Hand Hygiene

Hand hygiene is the most important measure to prevent the spread of infections. In the school setting hand hygiene is an important infection prevention method as a matter of habit with restroom use and food preparation. In the context of BBP and exposure control, hand hygiene should be enforced each time a staff member has an interaction with a student for standard first aid, medication administration, or direct care.

Hands should be washed prior to donning gloves, and after care is completed and gloves are removed.

Personal Protective Equipment (PPE)

PPE refers to wearable equipment that is designed to protect staff from exposure to or contact with infectious agents. PPE that is appropriate for various types of interactions and effectively covers personal clothing and skin likely to be soiled with blood, body fluid, or other OPIM should be available to school staff, in the appropriate sizes. These include gloves, face masks, protective eyewear, face shields, CPR masks, Kevlar sleeves or bite guards, and protective clothing (e.g., reusable or disposable gown). Contaminated PPE must be removed and disposed of before further contact with other surfaces or individuals occurs.

Examples of appropriate use of PPE for adherence to Standard Precautions include:

- Use of gloves in **all situations involving possible contact with blood or body fluids**, mucous membranes, non-intact skin (e.g., exposed skin that is chapped, abraded, or with dermatitis), or OPIM. Gloves must be removed and replaced as soon as practical when they are contaminated, torn, punctured, or when their ability to function as a barrier is compromised. Always wash or sanitize hands before donning gloves and after glove removal.
- Use of gowns to protect skin and clothing during procedures or activities where contact with blood or body fluids is anticipated, such as diapering, toileting, feeding, suctioning, general cleansing, or providing first aid.

- Use of mouth, nose, and eye protection (such as a face shield) during activities that are likely to generate splashes or sprays of blood or other body fluids, such as:
 - Feeding, providing care to, or assisting with a child with forceful vomiting or coughing
 - Suctioning a child with a tracheostomy with a history of forceful coughing or copious secretions
 - Assisting in the care of a student with a severe injury and spurting blood
 - Assisting a student with a head or facial wound
 - Assisting a student who displays erratic behavior that places the employee at risk (i.e. fighting, spitting).
- Use of Kevlar-sleeves or bite-guards when working with students with a history of biting or scratching behaviors.
- Use of mask, or face shield when respiratory transmission is of concern (see pandemic plan).

General Principles of PPE:

IF...	THEN...
It's wet (it's infectious)	Wear gloves
It could splash into your face	Wear a face shield
It's spread through the air	Mask yourself and the student
It could splash on your clothes	Wear a gown
You are providing direct care or first aid	Wear gloves
You are providing CPR	Use a barrier/CPR mask and gloves
There is a blood spill or body fluid spill	Summon BBP-trained staff for appropriate disinfection

Refer to Donning and Doffing (Applying and Removing) PPE addendum.

Respiratory Hygiene/Cough Etiquette

In the school setting, respiratory etiquette is an important measure to teach to staff and students as developmentally appropriate. Visual alerts, such as [Cover Your Cough](#) signage, can be used as a reminder for students and staff.

Appropriate respiratory etiquette includes practices on:

- Covering mouth and nose with a tissue when coughing or sneezing
- Using the nearest waste receptacle to dispose of the tissue after use
- Performing hand hygiene (e.g., hand washing with soap and water, alcohol-based hand sanitizer, or antiseptic hand wash) after having contact with respiratory secretions or contaminated objects/materials, such as a used tissue
- Sneezing or coughing into the elbow joint when a tissue and hand hygiene is not immediately accessible
- Avoiding sneezing directly into the hands, where they may subsequently contaminate other items or surfaces
- Having symptomatic individuals placed in a designated isolation space where risks to others are minimized until dismissed to home. A face covering can be offered if tolerated. Spatial separation of the person with a respiratory infection from others is important since respiratory droplets travel through the air for an average of 3-6 feet

To ensure these practices can be followed, each school should maintain the availability of materials for adhering to Respiratory Hygiene/Cough Etiquette in shared areas by:

- Providing tissues and no-touch trash receptacles for used-tissue disposal
- Providing conveniently located dispensers of alcohol-based hand sanitizer and, where sinks are available, ensure that supplies for hand washing (i.e., soap, disposable towels) are consistently available

Engineering and Work Practice Controls

Regulated Biohazardous Waste Containers

OR-OSHA defines regulated waste as “liquid or semi-liquid blood or other potentially infectious materials.” All school settings should be equipped with a regulated waste container to dispose of materials that are saturated with body fluids. There will be few items that are true regulated waste in a school setting, with the exception of contaminated sharps. Items with blood or other potentially infectious materials in a liquid or semi-liquid state, that if compressed would release liquid, must be disposed of in regulated waste containers that have a biohazard label. Contaminated items that are **exempt** from regulated waste include: diapers soaked with urine or feces and feminine hygiene napkins. The fluids are absorbed and contained easily within the product.

Determining what constitutes regulated waste (that must go into a biohazard-labeled container), versus contaminated waste (that can be disposed of in the regular trash) requires some independent decision and judgment. If waste soiled with blood or body fluids, as identified in the OR-OSHA definition of other potentially infectious materials, is saturated to the point of dripping, or would release fluids under compression, it needs to be identified, handled, and discarded as **regulated waste**. It is recommended that extra absorbent materials be utilized to clean up large body fluid spills, so that the waste is **not** liquid or dripping, and can therefore be disposed of in the normal trash as contaminated waste. School districts are encouraged to also review Oregon Statutes and Rules that govern the disposal of infectious waste.



BIOHAZARD

Sharps Safety

Needle stick injuries (piercing the skin with a used needle) are a potential risk in any work environment where medications are delivered via syringe or compatible device, or where lancets are used. In the school setting this is most often associated with the care of students who have medical conditions that require injectable medications. It is preferred that students provide self-care for such high-risk tasks whenever feasible, however this is not always safe due to the capability of the student or for the administration of emergency medications. Staff must be appropriately trained by a nurse before using injection devices. Handling of sharp instruments is covered with designated staff in specific training relative to their job responsibilities.

Certain controls must be enforced in any situation where sharps are present to reduce the risk of needle sticks, including:

- Avoid using needles that must be taken apart or manipulated after use.
- **Do not recap, bend, or break** one-time-use needles or other sharp devices. Two-hand manipulation of sharps is prohibited.
 - For example, students needing glucose monitoring by use of a lancet device should be encouraged to remove the lancet from the device themselves. If the student is unable, the use of a tool, such as needle nose pliers or medical clamps is required to prevent hand contact with the contaminated sharp.
- Always dispose of used needles or lancets in a sharps container appropriately labeled with a biohazard sign immediately after use.
- Never reuse needles and understand that needles should only be used a single time.
- Participate in specific training related to injectable medications.
- ALWAYS use a needle-stick-prevention tool in cases where a multi-use injection pen needs to be recapped after use. DO NOT EVER hold a cap, vial, or other container with your fingers while bringing the needle or sharp object toward the cap, vial, or container.

Contaminated sharps must be stored in closed puncture-resistant containers (sharps containers) with appropriate biohazard label and fill-line designation. It is the responsibility of the district to ensure the provision of a sharps container for each school/location along with timely removal and replacement when containers become full. This is usually accomplished through a contract with the local refuse service provider.

Health Room, Isolation Space, and Classroom Safety

Eating, drinking, applying cosmetics or lip balm, and handling contact lenses are prohibited in potential exposure areas. Storage of food/beverages is prohibited in places where body fluids or other potentially infectious materials may come in contact with surfaces such as countertops, sinks, or refrigerators in health rooms. Health rooms, isolation spaces, and classrooms should strive to have furniture items and other surfaces that are made of materials that are easily disinfected, and move toward removing surfaces that cannot be easily cleaned between uses. Sinks used for first aid and health room purposes should **not** have dishes, cups, or utensils placed in them. If the only sink available in the health room area is the first aid sink, a separate clean water source should be used to medicate and hydrate students.

Cleaning and Disinfecting Environmental Surfaces

The cleanliness of the district facilities is the responsibility of facility and custodial services who have specific expertise in the appropriate formulations to use for specific circumstances. For this reason, any body fluid spills should be immediately referred to custodial services. Schools should determine and implement an appropriate written schedule and documentation log for cleaning and decontamination of high contact surfaces and areas that may be susceptible to contamination with blood or other potentially infectious material, such as the health room and isolation room.

In the event of a blood spill, blood spill kits should be accessible on campuses. If custodial services are not immediately available, then another trained staff member who has gone through the in-person MESD BBP class should be called to clean the spill up immediately. Appropriate PPE should always be used with any body fluid clean up. Any articles used to clean body fluid spills must be handled with gloved hands and disposed of in an appropriate receptacle. If an absorbent agent is used, sweepings must be disposed of in a similar manner. Brooms and dustpans must be cleaned with a disinfectant. The CDC recommends using a freshly-mixed (no more than 24 hours old), diluted-bleach-solution (1 part bleach to 9 parts water) to disinfect areas with body fluid spills. Keep students away from areas being disinfected, or use a disinfectant from the Environmental Protection Agency's (EPA) List N with asthma-safer ingredients. If a bleach solution is not available, the school may use an EPA-approved tuberculocidal agent instead.

Steps for cleaning up a body fluid spill:

1. Absorb the fluid spill
2. Wash the surface with a general cleaner
3. Ensure that all surfaces are visibly clean prior to using the disinfecting solution (or the solution may be rendered ineffective)
4. Disinfect the area with a 1:10 bleach solution (or other EPA approved solution)
5. Leave the disinfectant on the contaminated surface for 10 minutes for blood spills, or follow the manufacturer's directions.

Using these basic principles, the management of spills should be flexible enough to cope with different types of spills, taking into account the following factors:

- the type of spill – sputum, vomit, feces, urine, blood or laboratory items
- the pathogens most likely to be involved in these different types of spills – stool samples may contain viruses, bacteria or protozoan pathogens,
- the size of the spill – spot (few drops), small (<10 cm), or large (>10cm) circumference
- the type of surface – carpet or impervious flooring
- the location involved – that is, whether the spill occurs in a contained area (such as a science laboratory), common area or in a restroom
- whether there is any likelihood of bare skin contact with the soiled (contaminated) surface

All disposal of biohazard waste will be in accordance with [EPA guidelines](#). The directives for appropriate sanitizing and waste removal should come from the facilities department in coordination with a contracted, licensed refuse-disposal company.

Transmission-Based Precautions

- Contact Precautions
- Droplet Precautions
- Airborne Precautions

Transmission-Based Precautions are the second tier of basic infection control and are to be used in addition to Standard Precautions for individuals with certain symptoms and/or diagnoses to prevent the spread of infectious agents.

Contact Precautions

Contact Precautions are to be used when a student has a known or suspected illness that is easily transmitted by direct contact or by indirect contact with items in their environment. Illnesses requiring contact precautions may include, but are not limited to, open and draining lesions, stool incontinence, uncontrolled secretions, or a new generalized rash.

When a student presents with an illness that requires contact precautions the following steps should be taken

- **Ensure appropriate student placement:** the student should be removed from the classroom setting and placed in the symptom space while awaiting parent/guardian arrival.
- **Use personal protective equipment (PPE) appropriately:** if the student requires care, gloves must be worn. Unlike a clinical setting, it is unlikely that a gown will be needed as staff should not be providing wound care or procedures beyond basic first aid.
- **Limit student activity:** the student's activity should be limited to reduce additional opportunity for contamination of surfaces.
- **Prioritize cleaning and disinfection:** after the student has been dismissed to home, ensure the area the student was located during direct care is appropriately sanitized. If there is a risk of contamination in other settings such as the classroom, cafeteria, or playground, ensure areas are appropriately disinfected before use by another student.

Droplet Precautions

Use Droplet Precautions when students or staff have a known or suspected infection transmitted by respiratory droplets. Droplets are generated by an individual who is coughing, sneezing, or talking. In the school setting this may be more common during influenza season and during the circulation of novel viruses.

- **Source control:** includes putting a medical-grade surgical mask on any caregivers and on the ill individual.
- **Ensure appropriate student placement:** a student who becomes symptomatic when the risk of specific viruses is increased, should be placed in a room individually (if possible), with appropriate supervision.
- **Use PPE:** universal mask use is required in all healthcare settings, including the health room and symptom space. Additionally, staff should wear a mask any time they are providing care to or screening an individual who presents with respiratory symptoms, even if they are not in a designated healthcare setting.
- **Limit activity of the ill student:** the student or staff's activity should be restricted, outside of the symptoms space, except for travel as needed to dismiss to home.

Airborne Precautions

Use Airborne Precautions when students or staff have a known or suspected infection transmitted by the airborne route (e.g., measles, chickenpox). Airborne precautions will rarely be used in the school setting; however, it is important to use identified control measures as vaccine preventable respiratory diseases are on the rise related to increased vaccine hesitancy.

- **Source control:** includes putting a face covering/mask on the ill individual.
- **Ensure appropriate student placement:** the student should be placed in a room individually (if possible), with appropriate supervision. If a symptom space is not available, ensure the student is separated from other students and staff.
- **Use PPE:** including a fit-tested NIOSH-approved N95 or higher level respirator for individuals providing direct care or supervision of the student. If a fit-tested NIOSH-approved N95 is not available, then a medical-grade surgical mask should be worn.

- **Limit activity of ill student:** the students activity should be restricted, outside of the symptom space, except for travel, as needed, to dismiss to home.
- **Notify susceptible individuals and advise immunization:** following contact with an individual identified as having a vaccine preventable disease, individuals susceptible to any diagnosed infection, such as measles, pertussis or varicella, should be advised to immediately seek immunization against infection. It is important to note that the school district cannot compel anyone to immunize themselves or their student, but students and staff who are unvaccinated (or not up to date with their vaccines) can be excluded for the maximum incubation period of a vaccine-preventable disease, up to 21 days from their last exposure.

Body Fluid Exposure Incidents

A body fluid exposure incident is an event where the potential or risk of exposure to infectious disease has occurred. This can occur through a variety of ways, but in the school setting this primarily occurs through contact with body fluids via mucous membranes or open, unprotected skin; through a human or animal bite; or through a needle stick injury.

When an exposure has occurred, the affected staff should immediately clean the affected area and follow the MESD Body Fluid Exposure Protocol (below). Students who are exposed to the body fluids of others should follow the initial preventive steps of the Body Fluid Exposure Protocol. The building administrator should then call the parent/guardian and recommend they call the students healthcare provider with any concerns or questions. This should be followed up with a school district incident report completed by the staff member who witnessed or reported the exposure event.

Needle-Stick Injuries

If a student or staff's skin is pierced or punctured with an object that may be contaminated with the body fluid of others, immediate first aid should occur including:

- Wash the wound thoroughly with soap and running water.
- Dry the wound and cover it with a clean dressing.
- Immediately notify the building administrator and call the MESD for assessment/referral.
- As soon as feasible, complete a school district incident report.
- Staff may be referred immediately to an occupational health clinic for further assessment and possible treatment, and may be required to report back to the provider for subsequent blood tests.
- Students should be referred to their healthcare provider.

Mucous Membranes

Any potential body fluid exposure toward the face, including the nose, mouth, eyes, or skin should be immediately flushed with running water. Staff should follow the MESD's Body Fluid Exposure Protocol (below).

Bites

While blood borne pathogen transmission is less common via bites, concerns of other infectious diseases may be present. Staff should follow MESD's Exposure Protocol (below).

Dog bites should be reported to the Multnomah County Animal Control at (503) 988-7387 to assess for rabies.

MESD Body Fluid Exposure Protocol

For ANY accidental body fluid contact to unprotected skin or mucous membrane (eyes, nose, or mouth), or any punctures through the skin (including bites):

**KEEP
CALM
AND
ASK FOR
HELP**

Get another adult to immediately take over your responsibilities so that you can take care of yourself right away.



For **any** body fluid contact to skin, **IMMEDIATELY wash skin** with soap and water for 2 minutes. For body fluid contact with face, **flush eyes, nose, and mouth wash** for 10 minutes with running water (remove contacts first).



You must **contact the MESD** as soon as possible at (503) 257-1732 for:
all blood exposures
or any body fluid contact to eyes, nose, or mouth
or all human bites that break skin
or any puncture with a contaminated object
or if you have any questions.



Notify your supervisor - same time; same day.

Complete all required district forms (at minimum an incident report form) and give them to your supervisor or the nurse consultant as directed.

The MESD Nurse will perform a full assessment of all occupational exposures as required by OSHA law, and refer employees for immediate follow-up at an occupational health clinic as necessary. This follow-up appointment, and any resulting care, treatment, or future appointments resulting from an occupational exposure are to be paid for by the employing district. It is assumed that all employees will follow their training and protocol on body fluid exposures, and notify their district supervisor of all body fluid exposure incidents and whether or not they were referred by the MESD Nurse for follow-up care.

PANDEMIC PLAN

A pandemic occurs when an infectious disease has spread globally. Viruses, such as coronaviruses and influenza, are routinely surveyed due to the propensity for mutations, human to animal transmission, and potential for pandemic events. The purpose of this document is to provide guidance for Non-Pharmaceutical Interventions (NPIs) and their use during a novel viral respiratory pandemic. NPIs are actions, apart from getting vaccinated and taking medications, that people and communities can take to help slow the spread of respiratory illnesses such as pandemic flu or novel coronaviruses. NPI's, specifically in regards to pandemic planning, are control measures that are incrementally implemented based on the level of threat to a community. This document should be used as a contingency plan that is modified with a response planning team based on the current level and type of pandemic threat.

Seasonal Respiratory Illness and Seasonal Influenza

Seasonal Respiratory Illness

There are several viruses that routinely circulate in the community that cause upper respiratory illnesses. These viruses include rhinoviruses, coronaviruses, adenoviruses, enteroviruses, respiratory syncytial virus, human metapneumovirus, and parainfluenza. The “common cold” is caused by rhinoviruses, adenoviruses, and coronaviruses. The symptoms of these seasonal illnesses may vary in severity but include cough, low-grade fever and sore throat (SDDH, 2019; Weatherspoon, 2019).




Seasonal Influenza

Influenza (flu) is a contagious respiratory illness caused by influenza viruses. There are two main types of influenza (flu) virus: Types A and B. The influenza A and B viruses that routinely spread in people (human influenza viruses) are responsible for seasonal flu epidemics each year. Influenza can cause mild to severe illness. Serious outcomes of flu infection can result in hospitalization or death. Some people, such as older people, very young children, and people with underlying health conditions or weak immune systems, are at high risk of severe flu complications. Routine symptoms associated with flu include fever, cough, sore throat, runny nose, muscle aches, headaches, fatigue, and vomiting (CDC, 2020).

Novel, Zoonotic, and Pandemic Viruses

Novel viruses refer to those not previously identified. A novel virus may be a new strain or a strain that has not previously infected human hosts. When a virus that has historically infected animals begins to infect humans it is referred to as a zoonotic virus. Pandemic refers to an epidemic that has spread over a wide geographical area or worldwide. The most common viruses associated with novel and pandemic outbreaks are influenza A and variants of coronaviruses. A flu pandemic occurs when a new variant that is different from seasonal viruses emerges and spreads quickly between people, causing illness worldwide. Most people will lack immunity to these viruses. Pandemic flu can be severe, causing more deaths than seasonal flu. Because it is a new virus, a vaccine may not be available right away. A pandemic could, therefore, overwhelm normal operations in educational settings (CDC, 2016b).

Differences between seasonal flu and pandemic flu:

Seasonal Flu	Mild to Moderate Pandemic	Severe Pandemic
 <p>THE VIRUS</p> <ul style="list-style-type: none">Caused by influenza viruses that are closely related to viruses that have previously circulated; most people will have some immunity to it.Symptoms include fever, cough, runny nose, and muscle pain.Complications such as pneumonia are most common in the very young and very old and may result in death.Vaccine is produced each season to protect people from the three influenza strains predicted to be most likely to cause illness. <p>IMPACT ON THE COMMUNITY</p> <ul style="list-style-type: none">Seasonal flu kills about 36,000 Americans each year and hospitalizes more than 200,000 children and adults.	 <p>THE VIRUS</p> <ul style="list-style-type: none">Caused by a new influenza virus that has not previously circulated among people and that can be easily spread.Because most people will have no immunity to the new virus, it will likely cause illness in high numbers of people and more severe illness and deaths than seasonal influenza.Symptoms are similar to seasonal flu, but may be more severe and have more frequent serious complications.Healthy adults may be at increased risk for serious complications. <p>IMPACT ON THE COMMUNITY</p> <ul style="list-style-type: none">May cause a moderate impact on society (e.g., some short-term school closings, encouragement of people who are sick to stay home).	 <p>THE VIRUS</p> <ul style="list-style-type: none">A severe strain causes more severe illness, results in greater loss of life, and has a greater impact on society.During the peak of a severe pandemic, workplace absenteeism could reach up to 40% due to people being ill themselves or caring for family members. <p>IMPACT ON THE COMMUNITY</p> <ul style="list-style-type: none">Schools and day care/child care facilities may be closed.Public and social gatherings will be discouraged.The patterns of daily life could be changed for some time with basic services and access to supplies possibly disrupted.

(Image: CDC)

Control Measures

While prophylactic vaccines and medications are appropriate interventions in some viral respiratory conditions, such as seasonal influenza, these are not always accessible for novel strains. NPIs are essential actions that can aid in the reduction of disease transmission. It is important to note that disease that is widely spread in the community has many options for transmission beyond the school setting, and the school district can only account for NPI's in the school setting and at school-sponsored events (CDC, 2017).



Personal NPIs are everyday preventive actions that can help keep people from getting and/or spreading flu. These actions include staying home when you are sick, covering your coughs and sneezes with a tissue, and washing your hands often with soap and water.



Community NPIs are strategies that organizations and community leaders can use to help limit face-to-face contact. These strategies may include increasing space between students in classrooms, making attendance and sick-leave policies more flexible, canceling large school events, and temporarily dismissing schools.



Environmental NPIs are surface cleaning measures that remove germs from frequently touched surfaces and objects.

(image: CDC)

Everyday Preventive Measures

Control measures to limit the spread of communicable diseases should be an active part of the school comprehensive and preventive health services plan. Routine control measures include:

- Hand hygiene (washing your hands for 20 seconds with soap and water with appropriate friction).
- Respiratory etiquette (cover your coughs and sneezes with a tissue and then throw it in the garbage after each use, or cough or sneeze into your elbow)
- Routine disinfection of shared items and frequently touched surfaces
- Staying home when sick, or experiencing any symptoms of illness

Control Measures for Novel or Variant Viruses

Control measures associated with novel or variant viruses are based on the severity and incidence of the specific virus. Some novel viruses are so mild, or rare, they may go undetected, while others are more contagious or severe. Since new viruses have no historical context, public health guidance evolves as case numbers increase and patterns are identified.

That being said, historical pandemic responses have provided a baseline set of evidence-based practices to create a framework for pandemic response planning in the school setting.


Control measures are incremental based on the current situation. The current situation will be defined by the public health officials based on the severity, incidence, and prevalence in the school setting. Level based responses will then be identified - these are generally defined as using a mild, moderate, and severe category, or for the purposes of this document, level 1, 2, and 3.

When cases of novel viruses are identified globally

When a novel disease is identified, it is the due diligence of school health services personnel and school administration to follow trends. When a novel strain is identified, routine prevention and exclusion measures should continue. In cases where student or staff travel is restricted secondary to pandemic events, it is the staff and parent/guardian's responsibility to communicate this restriction to the school district.

When cases of novel viruses are identified regionally or nationally

When a novel disease is identified in the U.S., it is important to identify the geographic spread and maintain awareness of specific public health messaging and direction. The Centers for Disease Control and Prevention (CDC) will have current national guidance. When novel viruses emerge in our state, the Oregon Health Authority (OHA) will provide regional guidance. OHA will have an alert for pandemic-specific content that can be subscribed to for updates. An individual within the district should subscribe to this alert to keep the district updated. If the region impacted is in Multnomah County, the Local Public Health Authority (LPHA) will provide school-centered communication and will potentially host conference calls. When cases are identified in the local region, the Multnomah County Health Department (MCHD) will work with the MESD CD Team, who will liaise with each component school district, and take part in a response team composed of School Health Services staff and district or individual school leaders, as appropriate. Response teams should be comprised of individuals who can fulfill roles with expertise in district policy and administration, clinical information, human resources, building-level management, risk management, and facilities to meet the general structure of Incident Command. The response team should hold regular meetings.



When public health has deemed a novel virus a pandemic threat, defer to the CDC in order to establish an emergency response framework with key stakeholders. During this time, preparedness planning will need to be initiated for the continuity of education in the event of school closure.

LEVEL ONE ACTIONS: PREVENTION FOCUSED (VIRUS DETECTED IN THE REGION)

Personal NPIs	Community NPIs	Environmental NPIs	Communication
<ul style="list-style-type: none"> • Increase routine hand hygiene. • Use alcohol-based hand sanitizer when hand washing is not an option. • Cover coughs/sneezes, throw away tissues at each use, wash your hands. • Stay home when ill for at least 24 hours after fever free without the use of fever-reducing medication. 	<ul style="list-style-type: none"> • Identify baseline absentee rates to determine if rates have increased uncharacteristically. • Increase communication and education on respiratory etiquette and hand hygiene in the classroom (teachers can provide age-appropriate education). • Communicable Disease surveillance - monitoring and reporting student and staff illness. • Increase space between students in the classroom. • Instruct students in small, stable groups as feasible. 	<ul style="list-style-type: none"> • Increase sanitizing of frequently touched surfaces and shared objects. • Develop prevention and post-exposure sanitizing strategies based on current recommendations. • Isolate students who become ill at school until parent/guardian can pick up. • Discourage the use of shared materials in the classroom. 	<ul style="list-style-type: none"> • Provide communications to families, students and staff based on current public health guidance. • Provide communication to immunocompromised student families to defer to their healthcare provider in regard to continued attendance.

When cases of novel viruses are identified in the community or incidence is increasing.

When novel viruses are identified in the community, but not in students or staff, the district will defer to local public health guidance. Public health guidance will likely occur when the overall incidence in the community is increasing (despite the proximity to the school). It is important to note that the school district can only apply controls around the school setting and school-sponsored events and activities. The school district cannot advise control measures around private clubs, organizations, or faith based communities. Each of these congregate settings are responsible for following local public health guidance as well.

When local transmission is detected, planning for cancellation of events and potential for dismissal and academic-continuity should be prioritized. Plans for potential prolonged staff absences should be prioritized, as well.

LEVEL TWO ACTIONS (IN ADDITION TO LEVEL ONE ACTIONS): INTERVENTION FOCUSED

Personal NPIs	Community NPIs	Environmental NPIs	Communication
<ul style="list-style-type: none"> Public health-specific guidance Be prepared to stay home if you or someone in your house is sick. 	<ul style="list-style-type: none"> Follow public health guidance Increase space between people at school following public health guidelines, as much as possible. Consider temporary dismissal of students attending childcare facilities and K-12 schools. 	<ul style="list-style-type: none"> Follow public health-specific guidance. Modify, postpone, or cancel large school events as coordinated with, or advised by, public health officials 	<ul style="list-style-type: none"> Work with LPHA to provide timely communication to families, students and staff about specific exposures. Provide communication to staff about the use of sick time and a reminder to stay home when sick. Advise parents to report actual symptoms when calling students in sick, as part of communicable disease surveillance.

When cases of novel viruses are identified in the school setting

When novel viruses are identified in the school setting, and the incidence is low, the LPHA will provide a direct report to the MESD CD Team on the diagnosed case. Likewise, the LPHA may impose restrictions on identified close contacts. However, it is important to note that if the incidence is high, the LPHA may not have the capacity to contact-trace and impose individual restrictions. As a result, the LPHA may create public statements that the school district should reiterate. If the school receives a report of a confirmed case, they should immediately contact the MESD CD Team who will work with the school nurse and building administrator to gather any additional information that will assist the health department with contact tracing efforts and advice isolation and quarantine, per established protocols.

LEVEL THREE ACTIONS (IN ADDITION TO LEVEL ONE & TWO): RESPONSE FOCUSED

Personal NPIs	Community NPIs	Environmental NPIs	Communication
<ul style="list-style-type: none"> Follow public health or government direction. 	<ul style="list-style-type: none"> Follow exclusion guidance designated by the LPHA, which may include physical distancing, revised gathering requirements, or school dismissal. 	<ul style="list-style-type: none"> Follow local public health direction on environmental cleaning. 	<ul style="list-style-type: none"> Coordinate with the LPHA to create communications to families, students and staff. Identify potentially impacted student populations, such as seniors and graduation track

POST EVENT

Personal NPIs	Community NPIs	Environmental NPIs	Communication
<ul style="list-style-type: none">• Routine hand hygiene and respiratory etiquette when LPHA deems processes may return to baseline.• Stay home when ill and until 24 hours fever free without the use of fever-reducing medications and symptoms are improving.	<ul style="list-style-type: none">• Routine illness exclusion when LPHA deems processes may return to baseline.	<ul style="list-style-type: none">• Routine sanitizing when LPHA deems processes may return to baseline.	<ul style="list-style-type: none">• Routine illness prevention and exclusion communication.• Participate in post-event evaluation to determine what worked in a response plan and what needs to be revised.

Special Considerations

Employee Sick Leave

Administration and human resources should work together with their legal advisor to determine the need to temporarily revise or flex sick leave to accommodate any public health guidance in regard to lost work. Prolonged exclusion may occur with individuals who are close contacts to a confirmed case, who are identified as suspect cases or who suffer from severe or prolonged illness.

School Closures

If school closure is ordered by the state, the district will abide the executive order. If a closure is advised by the LPHA, consultation should occur between legal, union, and district administration to ensure processes are consistent with [legal preparedness processes](#).

Immunocompromised Students

Students with immunocompromising health conditions and treatments may require exclusion from school outside of public health guidance. These students should provide documentation from their provider. The change in placement should be accommodated, as appropriate, under the Individuals with Disabilities Education Act (IDEA) and Free Appropriate Public Education (FAPE).

GLOSSARY & REFERENCES

Glossary of Terms

Administrative controls: Measures used in conjunction with engineering controls that eliminate or reduce a hazard by following established safe work practices and procedures for accomplishing a task safely.

Airborne precautions: Precautions that are utilized to protect against airborne transmission of infectious agents. Diseases requiring airborne precautions include, but are not limited to: Measles, Severe Acute Respiratory Syndrome (SARS), Varicella (chickenpox), and Mycobacterium tuberculosis.

Antibody: A protein produced as an immune response against a specific antigen.

Antigen: A substance that produces an immune response.

Bacteria: Microscopic living organisms. Some bacteria are beneficial, some are harmless, and some can be pathogenic (cause disease).

Biohazard Label: Legend (usually red or orange in color) to identify blood, regulated waste, or other potentially infectious materials (OPIM).

Biological Hazard: Any viable infectious agent that presents a potential risk to human health.

Blood borne pathogens: Microorganisms that can cause diseases, such as human immunodeficiency virus (HIV), hepatitis c virus (HCV), and hepatitis B virus (HBV), which are spread through contact with infected blood.

Cohort: A stable group of individuals who spend an extended period of their day in the same space together. In schools, this is likely a classroom, athletic team, before or after school group, bus route, etc. To reduce potential spread of communicable diseases in schools, cohorts should be kept as small as feasible

Communicable Disease: Illness that spreads from one person to another through contact with the infected person or their bodily fluids, or through contaminated food/water or disease vectors, such as mosquitos or rodents.

Contact Tracing: Public health practice that identifies and notifies individuals who have been exposed to an infectious disease.

Contaminated Waste: Items or surfaces with the presence or reasonably-anticipated presence of blood or OPIMs. These are items that are contaminated with body fluids, but with the capacity to absorb and contain those fluids so that they will not pool or drip. Examples of contaminated waste are diapers, sanitary pads, bandages, minor dressings from wounds or first aid treatment, and PPE. Contaminated items do not need to be placed in a Regulated Waste Container unless they are not able to contain the fluid and there is the possibility that they will allow dripping or pooling of the fluid within the container.

Decontamination: The use of physical or chemical means to remove, inactivate, or destroy the bloodborne pathogens on the surface or item to the point where they are no longer capable of transmitting infectious particles. The surface or item is rendered safe for handling, use or disposal.

Droplet precautions: Preventive methods used for diseases or germs that are spread in tiny droplets caused by coughing and sneezing (examples: pneumonia, influenza, whooping cough, bacterial meningitis).

Engineering Controls: Adjustments to the environment or to processes that eliminate or reduce exposure to hazard (for example, improving the ventilation system in a building).

Epidemic: A disease affecting a large number of people in a community or region.

Exclusion: Preventing someone from entering a place or participating in an activity.

Exposure Incident: A mouth, eye, nose, (mucous membrane), non-intact skin, or parenteral contact with blood or other potentially infectious materials that results from the performance of an employee's duties.

Immunocompromised: Having a weakened immune system, which limits the body's ability to fight disease, and therefore cannot respond normally to an infectious agent.

Isolation: Separating symptomatic and infected individuals from the healthy population. A method of controlling the spread of a disease.

Medical Wastes/Infectious Wastes: Blood, blood products, bodily fluids, any waste from human and animal tissues; tissue and cell cultures; human or animal body parts.

Novel: New or previously unidentified, as in, novel coronavirus.

Other Potentially Infectious Materials (OPIM): Human bodily fluid or tissue (other than blood) that can harbor or spread bloodborne pathogens, including but not limited to: saliva, cerebrospinal fluid, semen, vaginal secretions.

Pandemic: An epidemic that spreads across countries or continents.

Parenteral: Piercing the skin-barrier or mucous membranes through such events as needle sticks, human bites, cuts, and abrasions.

Pathogen: A microorganism that can cause disease.

Personal Protective Equipment or PPE: Items used when exposure to hazards cannot be engineered completely out of normal operations and when safe work practices and administrative controls cannot provide sufficient additional protection from exposure to infectious or hazardous conditions. PPE includes such items as: gloves, gowns, face shields, bite guards, and masks.

Quarantine: Separates and restricts the movement of people who were exposed to a contagious disease, to monitor for any symptoms of illness.

Regulated Waste: Liquid or semi-liquid blood or other potentially infectious materials; contaminated items that would release blood or other potentially infectious materials in a liquid or semi-liquid state if compressed; items that are caked with dried blood or other potentially infectious materials and are capable of releasing these materials during handling, including contaminated sharps; and pathological and microbial wastes containing blood or other potentially infectious materials.

Restrictable Diseases: Diseases that require exclusion from work, school and/or childcare facilities for the protection of public health.

Sanitize: Reduce contaminants (viruses, bacteria) on an object or surface.

Seasonal Illness: Illnesses whose occurrence appears to be associated with environmental factors (temperature and humidity changes). For example, colds, and other upper respiratory illnesses are more common during the winter months when people are indoors more often.

Sharps: Any devices that can be used to cut or puncture skin. Examples include: needles, syringes, and lancets (used for checking blood sugar). Sharps must be disposed of in an approved container, to avoid bloodborne pathogen exposure.

Standard Precautions: A set of infection control practices used to prevent transmission of diseases that can be acquired by contact with blood, body fluids, non-intact skin (including rashes), and mucous membranes. These measures are to be used when providing care to all individuals, whether or not they appear infectious or symptomatic.

Surveillance: Collecting and analyzing data related to a disease in order to implement and evaluate control measures.

Transmission: How a disease spreads. There are four modes of transmission:

- Direct—physical contact with infected host or vector
- Indirect—contact with infected fluids or tissues
- Droplet—contact with respiratory particles sprayed into the air (sneezed or coughed)
- Airborne—dried droplets that can remain suspended in the air for long periods of time (e.g., tuberculosis)

Universal Precautions: Preventing exposure to blood borne pathogens by assuming all blood and bodily fluids to be potentially infectious, and taking appropriate protective measures.

Vaccine: A preparation that is used to stimulate the immune system to produce antibodies to reduce the likelihood of contracting the disease and/or suffer severe disease, including that which results in hospitalization or death.

Vaccinated, fully: The status of having received all recommended vaccine doses in a series at the appropriate age and/or intervals for a specific disease.

Variant: A difference in the DNA sequence, a mutation. Viruses can change and mutate, and these variant forms can be intractable to established vaccines and treatments

Vector: A carrier of a pathogen that can transmit the pathogen to a living host. Mosquitoes, fleas, ticks, and rodents are examples of vectors.

Work practice controls: Measures intended to reduce the likelihood of exposure by changing the way a task is performed. They include appropriate procedures for handwashing, sharps disposal, lab specimen handling, laundry handling, contaminated material cleaning (OSHA, 2019b), and prohibiting two-handed manipulation of contaminated sharps.

REFERENCES

Alameda County Public Health Department (2013) Communicable Disease. Retrieved from <http://www.acphd.org/communicable-disease.aspx>

ASCD (Association for Supervisors and Curriculum Development) (2020). WSCC. Retrieved from <http://www.ascd.org/programs/learning-and-health/wsc-model.aspx>

BC Center for Disease Control (BCDC) (2009) A quick Guide to Common Childhood Diseases. Retrieved from http://www.bccdc.ca/schools/Documents/EN_Guide_Childhood_Diseases.pdf

Centers for Disease Control and Prevention. (2020). Influenza. Retrieved from <https://www.cdc.gov/flu/about/index.html>

CDC (2020) When and how to wash your hands. <https://www.cdc.gov/handwashing/when-how-handwashing.html>

CDC (2016a) Bloodborne Pathogens. Retrieved <https://www.cdc.gov/niosh/topics/bbp/emergnedl.html>

CDC (2016b) Variant Influenza Viruses: Background and CDC Risk Assessment and Reporting. Retrieved from <https://www.cdc.gov/flu/swineflu/variant.htm>

CDC (2017) Getting your school ready for pandemic flu. Retrieved from <https://www.cdc.gov/nonpharmaceutical-interventions/pdf/gr-pan-flu-ed-set.pdf>

Minnesota Department of Health (2020). Teaching Hand Hygiene. Retrieved from <https://www.health.state.mn.us/people/handhygiene/curricula/index.html>

Montana Department of Public Health and Human Services (MDPHHS) (2018) Communicable Disease: A guide for Schools in Montana. Retrieved from <https://opi.mt.gov/LinkClick.aspx?fileticket=Fy7meM6PXU%3D&portalid=182>

National Resource Center for Health and Safety in Child Care and Early Education (NRC), (2020). Environmental Health in Early Care and Education. Retrieved from https://nrckids.org/CFOC/Environmental_Health

Oregon Department of Education (2022) Communicable Disease Guidance for Schools. Retrieved from [Oregon Department of Education Communicable Disease Guidance Document](https://www.oregon.gov/education/communicable-disease-guidance-document).

South Dakota Department of Health (2019) Seasonal Respiratory Viruses. Retrieved from <https://doh.sd.gov/diseases/infectious/diseasefacts/viral-respiratory.aspx>

Victoria State Government. (2020). Managing spills of blood and body fluids and substances. Retrieved from <https://www.health.vic.gov.au/infectious-diseases/managing-spills-of-blood-and-body-fluids-and-substances>

Virginia Department of Health (2011) FAQ Respiratory Hygiene and Cough Etiquette. Retrieved from https://www.vdh.virginia.gov/content/uploads/sites/3/2016/01/RespiratoryHygieneCoughEtiquette_FAQ.pdf

Weatherspoon, D. (2019) Acute Viral Respiratory Infections. Retrieved from <https://www.healthline.com/health/acute-respiratory-disease>

COVID-19 SPECIFIC COMMUNICABLE DISEASE MANAGEMENT ADDENDUM

This plan is intended to be used in conjunction with the districts School Health Services Manual, School-Level COVID-19 Management Plan, Communicable Disease Plan, Pandemic Plan, and Exposure Control Plan, to meet the requirements of COVID-19 specific interventions in the school setting as designated by the Oregon Department of Education. This document addresses district specific processes to comply with the listed interventions. This document also uses guidance from the CDC's [Guidance for COVID-19 Prevention in K-12 Schools](#).

[Oregon Department of Education](#)

[Oregon Health Authority](#)

[Local Public Health Authority](#)

[Centers for Disease Control & Prevention](#)

Important Contacts

MESD Communicable Disease Team – cdsupport@mesd.k12.or.us

[MESD Coordinator of School Health Services](#) – 503-257-1732

[Multnomah County Health Department Communicable Disease Team](#) – 503-988-3406

Applicable Legislation

Emergency Rules Related to COVID-19

The Oregon Health Authority (OHA), Public Health Division, has temporarily adopted [OAR 333-017-0800](#) and OAR 333-018-0000 which adds a definition of COVID-19 and adds COVID-19 to the list of diseases reportable to public health authorities within 24 hours.

In addition, OHA is also adopting [OAR 333-019-0010](#) related to exclusion from schools, children's facilities, food service facilities and health care facilities

Existing Rules and Statutes

School Centered

OAR 581-022-0107 Operational Plans must be submitted for the 2022-23 school year

[OAR 581-022-2220](#) Standards for Public Elementary and Secondary Schools: Health Services

[OAR 581-022-2225](#) Emergency Plan and Safety Programs

[OAR 166-400-0010](#) Educational Service Districts, School Districts, and Individual School Records

[OAR 333-019-1030](#) Investigation and Control of Diseases: General Powers and Responsibilities

[ORS 433.255¹](#) Persons with or exposed to restrictable disease excluded from school or children's facility

[ORS 336.201¹](#) Nursing services provided by district

Occupational Health Centered

[1910-1030](#) OSHA Bloodborne Pathogens

Public Health Centered

[OAR 333-003-0050](#) Impending Public Health Crisis: Access to Individually Identifiable Health Information

[ORS 431A.015¹](#) Authority of Public Health Director to take public health actions

Recommended Staff Training and Communication

Designated Personnel

Designating staff for specific roles associated with COVID-19 prevention is important to ensure appropriate control measures are observed in a consistent manner and to ensure that data collection is accurate and appropriate for communicable disease response. Individual school buildings are required to outline roles and responsibilities related to communicable disease response. Designated staff must be identified within the School-Level COVID-19 Management Plan.

Staff Communication

- It is recommended that all staff receive communication on when to stay home and when they will be required to leave work based on illness.
- All staff will receive information on when to return to school based on ODE/OHA Communicable Disease Guidance for Schools and/or public health guidance.
- It is recommended that staff be informed of isolation/symptom space location and process for referral.
- It is recommended that staff be informed on infection control procedures.
- Formal communications to staff may will occur electronically.

General Staff Training

- All staff should be trained and advised on the logistical, operational, and physical changes in each building to maintain infection control and appropriate operational practices associated with cohorting or physical distancing.
- All staff who interact with students should be trained on passive screening and identification of excludable symptoms to determine when a student should be referred for full symptom screening and isolation.
- Staff will be trained on appropriate use, care, application, and removal of PPE.

Background Information

COVID-19 is an infection caused by a new coronavirus. Coronaviruses are a group of viruses that can cause a range of symptoms. While many cause mild illness, some, like COVID-19, can also cause more severe symptoms. COVID-19 infection often causes fever, cough, and trouble breathing. COVID-19 has additionally been reported to cause symptoms such as muscle pain, sore throat, headache, fatigue, nausea, vomiting, diarrhea, congestion/runny nose, and loss of taste or smell. Some people with the virus have mild symptoms or no symptoms, while other people can get quite sick, and rarely people die from COVID-19-related complications (OHA, 2020).

COVID-19 is spread when people breathe in (or touch and expose themselves to) the droplets created when ill people cough, sneeze, sing, or talk. This can happen when someone is close to a sick person, within six feet. Rarely, people might catch COVID-19 by touching a surface that a person with the infection coughed or sneezed on, and then touching their own mouth, nose, or eyes. Coronaviruses cannot survive for long on surfaces, though, so this is not currently thought to be a common means of spread (OHA, 2020).

Executive orders to close schools and other public spaces in Oregon and across the globe have evolved to include the slow and incremental reopening public spaces. Relative to school districts, this requires coordinated infection control planning for the upcoming school year with a framework for specified areas of intervention for reopening.

Pediatric Populations

COVID-19 is an infection caused by a novel coronavirus (SARS-CoV-2). Coronaviruses are a group of viruses that can cause respiratory symptoms ranging from mild to severe (CDC, 2020). Mild illness typically resolves without medical care. However, some variants, including COVID-19, can cause more severe or fatal diseases. COVID-19 infection often causes typical respiratory symptoms such as: fever, cough, and trouble breathing. COVID-19 has additionally been reported to cause symptoms such as muscle pain, sore throat, headache, fatigue, nausea, vomiting, diarrhea, congestion/runny nose, and new loss of taste or smell. Most people with a COVID-19 infection have mild symptoms or no symptoms, while other people, often the elderly or those who have underlying medical conditions, can get quite sick and possibly die from COVID-19 related complications (OHA, 2020).

COVID-19 is spread when an infected person breathes out droplets and very small particles that contain the virus. Those droplets and particles are then breathed in by other people or land on their eyes, noses or mouth. Rarely, people might catch COVID-19 by touching a surface that a person with the infection coughed or sneezed on, and then touching their own mouth, nose, or eyes. Anyone infected with COVID-19 can spread the virus even if they are not experiencing any symptoms of illness (OHA, 2020).

Pediatric Populations

It is important to note that complications from COVID-19 are well documented in elderly and fragile populations. Likewise, individuals testing positive for COVID-19 are more frequently recognized in older individuals with chronic health conditions. Current research shows that children most frequently resolve mild to moderate illness with symptomatic care only, and rarely encounter severe disease or critical complications (Hasan, A., Mehmood, & Fergie, 2020).

Although the risk of infection and complication is currently not thought to be high in student groups, students that have underlying medical conditions, are immunocompromised, or who live with someone who is at high-risk of disease must be accounted for. Accommodation for staff and students who are in high risk categories maybe necessary and should be driven by their primary health care provider. Consideration must be paid to the reviewing of 504 plans, Individualized Education Plans (IEP), and Individualized Health Plans (IHP) of vulnerable populations.

Vulnerable Populations

Ready Schools, Safe Learners identifies high risk populations at risk for severe illness from COVID-19 as people who have one or more of the following characteristics or conditions:

- » Age 65 years or older
- » Cancer
- » COPD chronic obstructive pulmonary disease
- » Serious heart condition
- » Immunocompromised state from solid organ transplant
- » Obesity
- » Type 2 diabetes mellitus
- » Chronic kidney disease
- » Sickle cell disease
- » Smoking
- » Pregnancy
- » People with disabilities and developmental disorders
- » Other conditions or risk factors identified by OHA, CDC, or a licensed healthcare provider

Students and staff with specific underlying conditions may be at increased risk of complications from COVID-19 and it may be necessary to provide specific accommodations for these individuals to ensure safety.

Families of students who are high risk may present healthcare provider orders indicating when they must stay home beyond that of a general student.

Due to the high volume of students with asthma, communication should be made via district and/or school newsletter to advise staff and families with students who have asthma should connect with their primary healthcare provider before school starts to determine any measures that may be necessary for that student individually. IHPs, 504s and IEPs may need to be revised by the education team to account for changes in provider orders, health restrictions, or to consider school accommodations

It is also important to remember that many students and staff members may have fragile family or household members, and accommodations may be requested for those situations.


Due to the nature of health privacy, staff members have the option to self-identify as high-risk and may provide recommendations from their healthcare provider, to district Human Resources. While they do not

have to provide a diagnosis to the district, documentation from the healthcare provider on restrictions may be necessary.

Communicable Disease Management

Existing Communicable Disease Management Plan should be deferred to for standards in disease control and prevention. Increased attention and education should be paid to these measures, including reinforcing routine measures such as hand and respiratory hygiene, establishing screening protocols and isolation and exclusion practices, and ensuring access to appropriate PPE. Individual school buildings are required to submit a School-Level COVID-19 Management Plan in which outbreak prevention, response and recovery must be addressed. Schools must also outline what mitigation measures will be utilized based on COVID-19 transmission within their facilities.

Layered Health and Safety Measures to Limit Spread of Disease



Cleaning and Disinfection

Routine sanitation measures should will be in full effect in the school building, including processes to respond to potentially infectious material as outlined in the Exposure Control Plan.

- All frequently touched surfaces (e.g., playground equipment, door handles, sink handles, drinking fountains) within the school and on school buses should will be cleaned and disinfected at least daily and between use as much as possible.
- Use of shared objects (e.g., gym or physical education equipment, art supplies, toys, games) should be limited when possible, or cleaned between use.
- It is recommended that students who are immunocompromised or in otherwise vulnerable populations for COVID-19 illness have their own supplies.
- A schedule should be designated by the Facilities Manager for increased routine cleaning and disinfection during pandemics.
- As necessary, additional custodial staff should will be deployed.
- Ensure safe and correct use and storage of cleaning and disinfection products, including storing products securely away from children. Use products that meet EPA disinfection criteria.
- Cleaning products should not be used near children, and staff should ensure that there is adequate ventilation when using these products to prevent children or themselves from inhaling toxic fumes.

Hand Hygiene and Respiratory Etiquette

There are some infection control measures that should be consistently practiced, taught, modeled, and reinforced in the school setting, even outside of pandemic spread. These primary principles include [hand hygiene and respiratory etiquette](#):

- Teach and reinforce [handwashing](#) with soap and water for at least 20 seconds and increase monitoring to ensure adherence among students and staff, prior to eating, after restroom use, upon entrance to school/class and before and after recess.
 - If soap and water are not readily available, hand sanitizer that contains at least 60% alcohol can be used (for staff and older children who can safely use hand sanitizer).
- Encourage staff and students to cover coughs and sneezes with a tissue. Used tissues should be thrown in the trash and hands washed immediately with soap and water for at least 20 seconds.
 - Students and staff may also be encouraged to cough into their elbow and away from other individuals when tissues and handwashing is not immediately accessible.

Airflow and Circulation

Ensure ventilation systems operate properly and increase circulation of outdoor air as much as possible, by opening windows and doors or using a fan to circulate outside air into the building. Do not open windows and doors if doing so poses a safety or health risk (e.g., risk of falling, triggering asthma symptoms) to children using the facility. Do not prop open an exterior door and leave it unmonitored. Consider the use of portable high-efficiency particulate air (HEPA) filters in rooms with limited ventilation and in symptom spaces.

COVID-19 Vaccination

Vaccination and testing are important strategies within the layered mitigation framework that schools operationalize to prevent viral transmission, and improve student and staff health and attendance. COVID-19 vaccination is available to everyone 6 months and older and significantly decreases the risk of serious illness, hospitalization and death from COVID-19. In Oregon, all school staff and volunteers are required to receive a primary vaccine series for COVID-19 or be granted a medical or religious exemption (OAR 333-019-1030). All students are strongly encouraged to receive COVID-19 vaccines, including booster doses. Vaccination is especially important for people who are immunocompromised or live or care for individuals who are immunocompromised (CDC, July 2022).

COVID-19 Testing

Individuals may spread COVID-19 as early as two days prior to symptom onset and some individuals have asymptomatic infections. Access to testing identifies positive cases of COVID-19 early and infectious individuals are isolated/excluded from school, thereby preventing viral transmission. All types of tests may be offered at schools, clinics, pharmacies, hospitals or performed in homes. For all types of tests, isolation recommendations and exclusion from school is determined by both the individual's symptoms and the results of the test using the Oregon Novel Coronavirus Disease 2019 Interim Investigative Guidelines and ODE Standard Exclusion Guidelines.

Schools may choose to offer diagnostic testing and/or exposure testing through the OHA program (OHA 3560 COVID-19 Testing in Oregon's K-12 Schools, May 2022).

- Diagnostic testing is performed when a student or staff develops symptoms at school or school-sponsored activities using a rapid antigen test.
- Exposure testing is performed on an asymptomatic individual within ten days after a known exposure to someone with COVID-19.

Screening testing may also be offered at regular intervals or prior to events or travel on participants regardless of vaccination status, presence of symptoms or known exposure. In Multnomah County K-12 Schools screening testing is an opt-in program, in which PCR tests are performed weekly.

Physical Distancing (Social Distancing/Spatial Distancing)

Physical or spatial distancing is the intentional physical distance placed between individuals to limit the likelihood of respiratory droplets reaching other individuals.

Room Capacity

- Consider physical distancing when determining room capacity.
- As needed, commons areas or alternate spaces maybe used to place student cohorts for instruction time in order to increase physical spacing between students.

Modified Layouts

- Excess furniture can be removed from classrooms to allow for increased spacing of desks.
- Desks or seating should be 3 feet apart (to the extent possible).

Staggered Scheduling

- Arrival/ drop-off times may be staggered by location and cohort. Arrival should be adjusted by building as determined to be necessary and feasible.
- Departure times may be staggered by location and cohort to the extent feasible to avoid crowding.
- Staggered and extended hallway passing should be endorsed to reduce hallway congestion and promote physical distancing.
- Stagger mealtimes, when feasible, to promote physical distancing while eating and to minimize time spent waiting in lines.

Instruction & Activities

- Outdoor spaces should be used as much as feasible.
- When distancing cannot be maintained, staff will be encouraged to wear additional PPE beyond their face coverings.

Communal Spaces

- Communal and shared spaces (such as the cafeteria) should [be cleaned and disinfected](#) between cohort use.

Cohorting (Stable Student Groups)

Cohorts help contain viral transmission, especially when there is minimal mixing of cohort members. In particular, the size of the cohort matters for risk management. Student cohorting limits the number of exposed people when a COVID-19 case is identified in the school and minimizes school-wide disruptions in student learning.

Student cohorting not only helps to quickly identify exposed people, it also minimizes disruptions to learning, because only the cohort members would be affected by response to the exposure, rather than the entire school. Maintaining small, stable cohorts can decrease the risk of closure to in-person instruction. It is recommended that cohorts be established with minimum numbers where feasible, understanding that fewer encounters and smaller numbers per cohort lends to reduced transmission of infectious disease. A smaller cohort size of 24-36 is recommended for public health and safety (OHA, 2020). **Accurate attendance must be maintained for proper cohort tracking and to provide accurate exposure response.**

It is important to note that cohorting may be more important in elementary settings where physical distancing is less likely to be maintained (AAP, 2020a). Cohorting should not be established based on the context of ability or disability. Cohorts should be established by grade levels in elementary settings and where feasible in upper grade levels.

Elementary

- Student and staff groupings should remain as static as possible by having the same group of children stay with the same staff throughout the school day as well as throughout the week.
- Interaction between cohort groups should be limited.
- Sanitizing of common areas should be performed between each cohort's use.
- Rather than students moving to common spaces for special courses, staff should present to each classroom.
 - Staff moving between cohorts must practice hand hygiene in between interactions and use appropriate logs to track their building activities.

- Maintain student cohorts throughout the day as feasible.
- Where stable cohorts are difficult to maintain, other mitigation practices should be re-emphasized distancing during activities and instruction.
- As much as feasible, student movement throughout the building should be limited.
- In cases where open doors and windows impact the operational settings of the ventilation system, facilities management should be consulted.

Masks & Personal Protective Equipment (PPE)

Face Coverings and Masks

Wearing a face covering is an effective way to reduce the spread of COVID-19 and other respiratory viruses. When local COVID-19 transmission is high it is recommended that all individuals wear a face covering indoors at school and on buses. During periods of low or moderate transmission individuals should consider wearing a face covering to reduce the spread of COVID-19 and other viruses (ODE, 2022). Requirements around face coverings and PPE may evolve to align with current state public health guidelines, new or revised mandates, and ODE’s requirements for the school setting. It should be clarified that face coverings are not synonymous with masks. Face coverings may include masks, cloth covers, or shields.

The OHA has adopted OAR 333-019-1011 which outlines masking requirements to control COVID-19 in healthcare settings. Healthcare providers and patients are required to mask when in a health care setting. A healthcare setting is defined as any place where health care, including physical, dental, or behavioral health care is delivered (OHA, 2022). In schools, this may include, but is not limited to: health room, symptom space, isolation space, mental or behavioral health care setting and other areas that are designated for providing health care. Nurses, school staff, students and visitors are required to wear a mask in areas of the school that meet the definitions of a healthcare setting.

PPE

PPE is specialized clothing or equipment used by staff in an occupational setting to reduce the risk of infection transmission or risk of chemical exposure. The district Exposure Control Plan (pages 15-25) should be consulted for necessary and appropriate use of PPE. PPE will be advised based on the interaction with students or the risk involved related to frequency and type of interaction, volume and duration of interaction and the developmental stages and health status of the individuals involved. For example, clinical staff or staff in isolation spaces may require a higher level of PPE than staff teaching in an academic space.

Personal Protective Equipment/Barriers		
INDIVIDUALS	REQUIRED PPE	RECOMMENDED PPE (in addition to required PPE)
Staff providing care in the health room or symptom/isolation space	Medical grade PPE per Transmission Based Precautions Mask	No additional recommendations
Acutely ill student in isolation	Mask (until parent picks up)	No additional recommendations
Caregivers performing aerosol generating procedures	Fit tested NIOSH approved N95, gloves, face shield, gown if splash likely	Fit tested NIOSH approved N95, gloves, face shield
Additional Guidance for face covering use: <ul style="list-style-type: none"> • Cloth face coverings must be laundered regularly • New disposable face covering must be used daily • Face shields are reusable and should not be shared between staff • Face coverings cannot be shared • Plastic face shields should be disinfected regularly 		

N95 Masks and Surgical Masks

N95 respirators and surgical masks are examples of personal protective equipment that are used to protect the wearer from airborne particles and from liquid contaminating the face.

- NIOSH-approved respirators, such as N95s, provide the highest level of protection when compared to medical grade face masks and cloth face coverings. Please note that N95 masks should be fit tested, and the user trained for appropriate use. In addition, N95 masks must form a seal to the face and be applied and removed correctly for effective use.
- KN95 masks are filtering facepiece respirators that provide 95% protection against particulates. KN95 masks are designed and tested to meet international standards and do not have to be fit tested.
- Surgical masks are appropriate for cases where direct face to face interactions will occur, in order to create a physical barrier of protection. If worn properly, a surgical mask is meant to help block large-particle droplets, splashes, sprays, or splatter that may contain germs (viruses and bacteria), keeping it from reaching your mouth and nose. Surgical masks may also help reduce exposure of your saliva and respiratory secretions to others.

Cloth Face Coverings









[Cloth face coverings](#) are meant to protect other people in case the wearer is unknowingly infected but does not have symptoms. Cloth face coverings are not surgical masks, respirators, or other medical personal protective equipment. The most effective cloth face coverings are multiple layers and fit snugly around the nose, cheeks and chin. It is recommended that information be provided to staff, students, and students' families on [proper use, removal, and washing of cloth face coverings](#).

Staying Home When Appropriate & Exclusion of Students and Staff

It is crucial that school staff and families understand when individuals must stay home. It is important for all staff to role model appropriate behaviors. Communication should be made regularly to advise families not to send children to school ill and to remind staff not to report to work ill. Students and staff should be excluded per ODE/OHA Communicable Disease Guidance for Schools if they present to the school building with any symptoms of illness that require exclusion (below).







When Should I Keep My Student Home?

The list below gives school instructions, not medical advice. Please contact your healthcare provider with health concerns. **During 2020-2021, anyone exposed to COVID-19 must stay home for 14 days.**

Symptoms of Illness	Student May Return to School After... *The list below tells the shortest time to stay home. A student may need to stay home longer for some illnesses.
 Fever: temperature of 100.4F (38C) or greater	Fever-free for 24 hours without taking fever-reducing medicine AND after a COVID-19 test is negative , OR 10 days if not tested.
 New cough illness OR New difficulty breathing	Symptom-free for 24 hours AND after a COVID-19 test is negative , OR 10 days if not tested. If diagnosed with pertussis (whooping cough), the student must take 5-days of prescribed antibiotics before returning.
 Headache with stiff neck or with fever	Symptom-free OR with orders from healthcare provider to school nurse.
 Diarrhea: 3 loose or watery stools in one day OR not able to control bowel movements	Symptom-free for 48 hours OR with orders from healthcare provider to school nurse.
 Vomiting: one or more episode that is unexplained	Symptom-free for 48 hours OR with orders from healthcare provider to school nurse.
 Skin rash or open sores	Symptom-free , which means rash is gone OR sores are dry or can be completely covered by a bandage; OR with orders from healthcare provider to school nurse.
 Jaundice: (new) yellow color in eyes or skin	After the school has orders from healthcare provider or local public health authority to school nurse.
 Red eyes with eye discharge: yellow or brown drainage from eyes	Symptom-free , which means redness and discharge are gone OR with orders from healthcare provider to school nurse.
Acting different without a reason: unusually sleepy, grumpy, or confused	Symptom-free , which means return to normal behavior OR with orders from healthcare provider to school nurse.
Major health event , like an illness lasting 2 or more weeks OR a hospital stay	After the school has orders from healthcare provider to school nurse.
Student's health condition requires more care than school staff can safely provide	After measures are in place for the student's safety.

PLEASE KEEP STUDENTS WITH SYMPTOMS OUT OF SCHOOL

This list is school instructions, not medical advice. Please contact your health care provider with health concerns.

SYMPTOMS OF ILLNESS	THE STUDENT MAY RETURN AFTER...
 Fever: temperature of 100.4°F (38°C) or greater	<p style="text-align: center;">*The list below tells the shortest time to stay home. A student may need to stay home longer for some illnesses.</p> <p>*Fever-free for 24 hours without taking fever-reducing medicine AND per guidance for primary COVID-19 symptoms.</p>
 New cough illness	<p>* Symptoms improving for 24 hours (no cough or cough is well-controlled) AND per guidance for primary COVID-19 symptoms.</p>
 New difficulty breathing	<p>* Symptoms improving for 24 hours (breathing comfortably) AND per guidance for primary COVID-19 symptoms. Urgent medical care may be needed.</p>
 Diarrhea: 3 loose or watery stools in a day OR not able to control bowel movements	<p>*Symptom-free for 48 hours OR with orders from doctor to school nurse.</p>
 Vomiting: one or more episode that is unexplained	<p>*Symptom-free for 48 hours OR with orders from doctor to school nurse.</p>
 Headache with stiff neck and fever	<p>*Symptom-free OR with orders from doctor to school nurse. Follow fever instructions above. Urgent medical care may be needed.</p>
Skin rash or open sores	<p>*Symptom free, which means rash is gone OR sores are dry or can be completely covered by a bandage OR with orders from doctor to school nurse.</p>
Red eyes with colored drainage	<p>*Symptom-free, which means redness and drainage are gone OR with orders from doctor to school nurse.</p>
Jaundice: new yellow color in eyes or skin	<p>*After the school has orders from doctor or local public health authority to school nurse.</p>
Acting differently without a reason: unusually sleepy, grumpy, or confused.	<p>*Symptom-free, which means return to normal behavior OR with orders from doctor to school nurse.</p>
Major health event , like an illness lasting 2 or more weeks OR a hospital stay, OR health condition requires more care than school staff can safely provide.	<p>*After the school has orders from doctor to school nurse AND after measures are in place for the student's safety. Please work with school staff to address special health-care needs so the student may attend safely.</p>



(Image: Oregon Health Authority)

Identifying, and Isolating Ill Students and Staff


Identification of ill students and staff is crucial in illness prevention in school buildings. All staff, students, and families should have education provided on symptoms, in order to self-identify when developmentally possible. Screening should happen through a multi-layered approach:

1. Parents are educated on and asked to screen their students each morning before sending them to school
2. Bus drivers may passively screen students getting onto bus and notify school if student appears symptomatic
3. Teachers and staff typically provide passive screening throughout school day for symptoms of exclusion – send symptomatic individuals to the health room for further screening and possible isolation
4. Additional screening is provided in the health room to determine if there is a need to isolate the individual.

Health Promotion, Prevention, and At-Home Screening

Parents will be provided a parent symptom screening algorithm or check list (see example below), and be advised to screen their students prior to sending them to school. Parents should be notified on circumstances in which students should not attend school and when children will be excluded from school. Families and staff will additionally be provided with COVID-19 symptom checkers to use as tools to determine follow up. School staff should not provide medical advice.

- [CDC Self-Checker](#)



Isolate Those Who Are Sick

- Each school must have a designated isolation space with designated personnel to supervise the space. PPE must be available for designated staff to use [Standard and Transmission-Based Precautions](#), as per the *District Exposure Control Plan*.
- The district *Communicable Disease Plan* should be referred to for isolation measures.
- COVID-19 specific isolation measures will be updated as state public health guidance is updated.

Isolation Measures

- Immediately separate students who are determined to have symptoms meeting exclusion criteria to isolation/symptom space. .
- Students awaiting pick up are within line of sight of staff in the designated isolation area.
- Staff should wear a mask and maintain physical distancing, and never leave a child unattended.
- While exercising caution to maintain safety is appropriate when working with children exhibiting symptoms, it is also critical that staff maintain sufficient composure and disposition so as not to unduly worry a student or family.
- If more than one student is in an isolation space, appropriate distance should be maintained, and appropriate barriers and privacy must be in place.
- Health checks will be conducted safely and respectfully, and in accordance with any applicable privacy laws and regulations.
- Staff will maintain student confidentiality.
- Reinforce appropriate exclusion action with parents/guardians and staff using [ODE/OHA's Communicable Disease Guidance for Schools](#)

Isolation/Symptom Spaces

An appropriate isolation space as described in the *Communicable Disease Plan* and consistent with state legislation, should be accessible in each building. The intent is to mitigate the risk of transmission from an ill individual to well individuals.

The isolation space should observe public health guidelines to the extent feasible to ensure each element of infection prevention is followed as per *Transmission Based Controls* and COVID-19 guidance correctly. The isolation space should be logistically accessible in proximity to the health room, but not in such close proximity that infection risk is a concern.

CDC guidelines should be visited with the following five requirements in mind:

1. Isolation space must be separate from routine health room
2. Isolation space must have appropriate ventilation
3. Students must be supervised while in isolation space
4. Staff must have appropriate PPE while in the isolation space
5. Appropriate physical distancing, barriers, and confidentiality must be maintained in the isolation space

CDC Guidelines for Isolation Rooms

Isolation Space	CDC Guidelines
Physical distance	It is strongly recommended that ill students are physically distanced while in the isolation space to the greatest extent possible.
Cleaning and sanitizing	After dismissal of ill student, close off areas used by a sick person and do not use these areas again until after cleaning and disinfecting . When disinfecting the isolation space, wear a medical grade face mask, a face shield, and gloves. Ensure safe and correct use and storage of cleaning and disinfection products, including storing products securely away from children.
Ventilation	Designated isolation space should have adequate ventilation, i.e. exterior windows and/or ventilation fans. Ensure fans do not re-circulate into air supply; vent to exterior or into non-communicating space (wall voids, attic).
Hand hygiene	Care providers should wash hands frequently and thoroughly before and after providing care, including after removal of gloves. Ensure isolation space has ready access to soap and water. Sink at the entryway is preferred. If soap and water is not accessible, use hand sanitizer with $\geq 60\%$ alcohol content and wash hands with soap and water as soon as possible.

Isolation Space	CDC Guidelines
Mask/other PPE	<p>Staff tending to symptomatic individuals should wear, at a minimum, a medical-grade face mask. Do not discard the face mask when exiting the isolation room care space unless it is soiled. Gloves should be worn if there is risk of exposure to blood and body fluids. A face shield should be worn if there is a risk of body fluids splashing towards the face.</p> <p>Any PPE used during care of a symptomatic individual should be properly removed and disposed of prior to exiting the isolation room, and hands washed after removing PPE.</p>
Student safety and well-being	<p>Adjust protocols to age and developmental abilities. Ensure line of sight; keep the ill student visible. To reduce fear, anxiety, or shame related to isolation, provide a clear explanation of procedures, including use of PPE and handwashing.</p>

Communication Systems

The district will implement and provide communications, including health promotion, communication of policies and restrictions, and communication regarding potential exposures or exclusions.

School Communication

Signs and Messages

- Post [signs](#) in highly visible locations (e.g., school entrances, restrooms) that promote everyday protective measures and describe how to stop the spread of germs (such as by [properly washing hands](#) and [properly wearing a cloth face covering](#) where applicable)
- Broadcast regular [announcements](#) on reducing the spread of COVID-19
- Messages may be included on websites, in newsletters, and on district social media


Direct Communication

- Health promotion material should be shared with students, staff and families
- In addition to posting exclusion criteria on web pages and in newsletters, families should be advised on policies related to sick students, potential home isolation criteria, and student exclusion criteria
- Families and staff should receive communication on any logistical changes for arrival and departure, physical distancing, schedule changes, and non-pharmaceutical interventions employed
- Age appropriate classroom curriculum will be used to encourage positive hygiene behaviors
- Families will be advised to report if their student:
 - Has symptoms of illness,
 - Has had a positive test for COVID-19.

Communication Regarding Confirmed Cases

For a complete overview of communication and response related to confirmed communicable disease cases, refer to the [MESD Communicable Disease Communication Protocol](#).

- District specific protocols and practices should be communicated by the superintendent
- Building specific protocols should be communicated by the building administrator
- The MESD CD Team will inform school nurse, principal, and designated district contacts of all confirmed cases of COVID-19 communicated by the LPHA
- Any parent, student, or staff reports of illness should be reported to the school



- The CD Team will work with the LPHA to provide district administration with a letter to share with students, families and staff (if notification is determined to be necessary)

Staff Self Identification

Staff should be advised to report to administration, or their direct supervisor, if they:

- Have excludable symptoms of illness
- Have symptoms of COVID-19

Have had a positive test for COVID-19 Sick staff members or students should not return until they have met criteria to discontinue home isolation.

Public Health Communication

- The MESD CD Team are the liaison with the LPHA and the Deputy Health Officer
- The MESD CD Team are subscribed to any COVID-19 updates via [Oregon Health Authority](#) that reports the incident of disease and provides routine updates by region (OHA may vary the updates that are provided to the public)

Nutrition Services

Food Service personnel should follow all existing mandates on health and hygiene and food safety. Any specific measures or intervention should be coordinated with the Facilities Manager and the Nutrition Manager. Additional measures should be endorsed in response to COVID-19, and any potential outbreaks, to improve infection control measures around food services.

- Children should wash/sanitize hands prior to eating.
- Children using school lunch services should be served individually plated meals.
- School lunch times should be staggered to maintain spatial distancing to the extent feasible. Shared spaces should be sanitized between use.
- Use of disposable food service items is promoted when feasible (e.g., utensils, dishes). If disposable items are not feasible or desirable, ensure that all non-disposable food service items are handled with gloves and washed with dish soap and hot water or in a dishwasher.
- Individuals should wash their hands after removing their gloves or after directly handling used food service items.

Transportation

Measures taken on transportation should follow the processes of school operations to the extent feasible. Coordination with transportation services and the facilities manager should be ongoing.

Bus Drivers

- Transport vehicles (e.g., buses) that are used by the school, require that drivers practice all safety actions and protocols as indicated for other staff (e.g., hand hygiene, cloth face coverings/face shields).
 - Bus drivers shall wear PPE as designated under PPE section.
- All frequently touched surfaces on school buses will be [cleaned and disinfected](#) at least daily and between use as much as possible.

Bus Screening

- Bus drivers shall passively/visually screen students as they enter the bus, and attempt to mark attendance and any existence of symptoms of illness on a daily bus log. In recognition of transportation and safety measures, and the priority of the district to maintain student safety in all areas, buses will not remain stationed in the roadway for prolonged periods of time to assess students. Measures will be taken to isolate students who become ill on bus routes and as soon as students arrive at school:

IF...	THEN...
Student is visibly ill upon entry into the bus	The bus driver should request the student remain at home, if age appropriate or parent present. If parent/guardian not present, see below. All efforts should be made to maintain dignity of student and family.
Student is visibly ill and parents/guardians are not present	Student should be seated close to the front, and as separate from other students as feasible. Student should be immediately isolated upon arrival to school - school staff should report to bus to retrieve student and take to isolation space. All efforts should be made to maintain privacy and dignity of student.
Student becomes ill on bus route	Bus driver should contact dispatch to notify appropriate school and parent/guardian. Student should be immediately isolated upon arrival at school - school staff should report to bus to retrieve student and take to isolation space. All efforts should be made to maintain privacy and dignity of student.
Student is in distress during bus route	Follow existing emergency transportation procedures to contact Emergency Medical Services (EMS).

Distancing and Staggering

Create distance between children on school buses (e.g., seat children one child per row, skip rows) if feasible. When students unload from bus, they should be directed to exit the bus one at a time.

Maintaining Healthy Operations

Schools may consider implementing several strategies to maintain healthy operations.

Regulatory Awareness

- Be aware of local or state regulatory agency policies related to group gatherings to determine if events should be held

Visitors and Volunteers

- Communication should be made to essential visitors to indicate that they cannot report to buildings if they have been sick or recently tested positive for COVID-19
- Consider restricting non-essential visitors
- Physical distancing is recommended as a layered mitigation measure to reduce the spread of COVID-19
- Visitors should be asked to wash hands or use hand sanitizer upon arrival
- When face coverings are not mandated by law or executive order, schools may recommend visitors wear face coverings while on site.

Gatherings and Field Trips

- When local COVID-19 transmission is high, pursue virtual group events, gatherings, or meetings, if possible, and promote social distancing between people if events are held in person.
- If possible, limit activities involving external groups or organizations as possible and under executive orders – especially with individuals who are not from the local geographic area (e.g., community, town, city, county)
- Pursue virtual activities and events in lieu of field trips, student assemblies, special performances, school-wide parent meetings, and spirit nights, as possible
- Pursue options to convene sporting events and participation in sports activities in ways that minimizes the risk of transmission of COVID-19 to players, families, coaches, and communities

Participation in Community Response Efforts

- Consider participating with local authorities in broader COVID-19 community response efforts (e.g., sitting on community response committees)

Sharing Facilities

- Ensure that any organizations that share or use the school facilities also follow these considerations

Continuity of Routine School Health Services

Ongoing school health services must be provided in tandem with COVID-19 specific interventions. School Health Services should continue operations per established MESD policies, procedures, School Health Services Manuals, and service contracts.

Special consideration should be paid to where care (such as diabetic care or medication administration) is provided for high risk students in vicinity to isolation rooms.

REFERENCES

Centers for Disease Control and Prevention (CDC) (2020a). Human Coronaviruses. Retrieved from <https://www.cdc.gov/coronavirus/general-information.html>

Centers for Disease Control and Prevention (CDC) (2021). Reopening Guidance for Cleaning and Disinfecting Public Spaces, Workplaces, Businesses, Schools, and Homes. Retrieved from <https://www.cdc.gov/coronavirus/2019-ncov/community/reopen-guidance.html>

Oregon Department of Education and Oregon Health Authority (OHA/ODE) (2022). Communicable Disease Guidance for Schools. Retrieve from <https://www.oregon.gov/ode/students-and-family/healthsafety/Documents/commndisease.pdf>

Hasan, A., Mehmood, N., & Fergie, J. (2020). Coronavirus Disease (COVID-19) and Pediatric Patients: A Review of Epidemiology, Symptomatology, Laboratory and Imaging Results to Guide the Development of a Management Algorithm. *Cureus*, 12(3), e7485. <https://doi.org/10.7759/cureus.7485>

North Carolina Department of Health and Human Services (2022) Strong Schools NC Public Health Toolkit. Retrieved <https://covid19.ncdhhs.gov/strong-schools-nc-public-health-toolkit>

Oregon Health Authority (OHA) (2022). COVID-19. Retrieved from <https://govstatus.egov.com/OR-OHA-COVID-19>

Oregon Department of Education (ODE) (2020). Ready Schools, Safe Learners. Retrieved from <https://www.oregon.gov/ode/students-and-family/healthsafety/Documents/Ready%20Schools%20Safe%20Learners%202020-21%20Guidance.pdf>


COVID-19 TOOLS

- ▶ Parent Daily Symptom Checker Flowchart
- ▶ MESD Daily Illness Self-Screening Checklist
- ▶ Symptom Screening Flowchart
- ▶ Communicable Disease Sheet (CDS)
- ▶ Isolation Procedures
- ▶ Donning & Doffing (Applying & Removing) PPE

Parent Daily Symptom Checker Flowchart

Can my child go to school today?

Ask the questions in the 3 red boxes EVERY MORNING
Then follow the arrows based on your answer





DAILY ILLNESS SELF-SCREENING CHECKLIST

Published: 2/2021 Updated: 3/2022

Employees and parents/guardians of students will be required to self-screen daily for symptoms of illness before entering district property or using district transportation (employees self-screen, parents/guardians screen their children). In addition, this Self-Screening Checklist should be used to determine whether there is a risk for potential illness.


If an employee or student has any of the following symptoms, or tested positive for COVID-19, the employee or the student should stay home, report or have parent/guardian report the absence and symptoms to the school or appropriate supervisor, and contact their healthcare provider as necessary.

COVID-19 Diagnosis
<input type="checkbox"/> Employee or student has tested positive for COVID-19
Primary Symptoms of COVID-19
<input type="checkbox"/> Fever (temperature of 100.4°F or higher)
<input type="checkbox"/> A new cough that has not been diagnosed or cleared by a healthcare provider
<input type="checkbox"/> Unexplained shortness of breath
<input type="checkbox"/> New loss of taste or smell
<input type="checkbox"/> Chills
<input type="checkbox"/> Difficulty breathing
Additional Excludable Symptoms of Illness
<input type="checkbox"/> Vomiting
<input type="checkbox"/> Diarrhea
<input type="checkbox"/> Undiagnosed rash or sores

*For information about exclusion see the Oregon Department of Education and the Oregon Health Authority guides: [Planning for COVID-19 Scenarios in Schools](#) and [ODE/OHA Communicable Disease Guidance for the School Setting](#)

Symptom Screening Flowchart

The list of symptoms associated with COVID-19 is exhaustive, but there are specific indicators for isolation and exclusion. At the beginning of the day and as the day progresses, any visual observation of symptoms under *visual screening indicators* or symptoms reported under *student complaint indicators* should be immediately referred to designated screening personnel. These designated personnel should be identified in each building and staff should be advised of the process to communicate with these designated staff. Referral to designated staff will be non-discriminatory (not based on appearance or hygiene), and be respectful in nature



Isolation Procedures

Isolation spaces should meet criteria previously listed, and should not have signs that violate dignity or health privacy. Isolation spaces must have designated staff to monitor students, and the designated staff must have access to appropriate PPE and, preferably, a sink for handwashing at the isolation space. Staff must use [Standard and Transmission-Based Precautions](#), as per the District *Exposure Control Plan* and The district *Communicable Disease Plan*.

Upon referral to designated staff, staff should screen ill students for excludable symptoms. Individuals who screen positive for the following by designated staff must be isolated:

- Fever and chills [Take temperature (should be under 100.4 °F)]
- New or worsening cough
- Shortness of breath or difficulty breathing not explained by an underlying condition or relieved with rescue medication
- New loss of taste or smell
- Nausea or vomiting
- Diarrhea
- New undiagnosed rash or sores

For students with Individual Health Plans (IHPs), these plans should be deferred to if symptoms might be associated with underlying health conditions. For specific symptoms associated with asthma or altered level of consciousness, defer to district standard procedures. Any concerning symptoms should be referred to the school nurse. **Any student in distress should have EMS (9-1-1) called immediately.**

1. When it is determined that a student has symptoms that are excludable, initiate contact to parent/guardian for pick up.
2. Take the student to the isolation space pending parent/guardian pickup in a safe and respectful manner.
3. Explain to student the process in developmentally appropriate terms, avoiding phrases such as “isolation.”
4. Student will be provided a face mask. Ensure that this is removed if the student falls asleep to maintain safety.
5. Staff should wear a medical grade mask upon entry to isolation space.
6. Remain calm and practice measures to maintain student privacy, confidentiality, and dignity to the highest extent feasible.
7. Staff should promote respiratory etiquette and physical distancing in the isolation space.
8. If more than one student is in an isolation space, appropriate distance and privacy should be maintained between students.
9. Do not leave students unattended.
10. Reinforce appropriate exclusion action with parent/guardian using ODE/OHA’s COVID-19 Exclusion Summary Guidance for K-12 Schools.
11. Ensure appropriate sanitizing between each use of space.

Communicable Disease Sheet (CDS)

Each school should have a way to track reports of communicable diseases in their population at any given time so that efficient follow up by the nurse and MESD CD Team can be completed. This Communicable Disease Sheet should be shared between front office staff taking phone calls from parents reporting student absences due communicable disease, such as COVID-19. The school nurse and the school administrator or their designee will have access to the information on the CDS. The nurse will review the sheet and follow up with students as needed.

Ainsworth Elementary CDS ☆ 📄 ☁

File Edit View Insert Format Data Tools Extensions Help [Last edit was seconds ago](#)

100% | \$ % .0 .00 123 | Default (Ari... | 10 | **B** *I* A | 🗑️ 📏 📐 📑 📊 📈 📉 📋 📌 📍 📎 📏 📐 📑 📊 📈 📉 📋 📌 📍 📎

	A	B	C	D	E	F	G	H
1	Student/Staff Member name:	Student ID Number:	Best Contact Number:	Reason for follow-up:	Date communicated with parent/guardian:	Time communicated with parent/guardian:	Nurse updates:	
2								
3								
4								
5								
6								
7								
8								
9								
10								
11								
12								

Donning & Doffing (Applying & Removing) PPE

Donning (applying) PPE

1. Perform appropriate hand hygiene.
2. [Determine appropriate PPE to be worn](#) depending on duties being performed.
3. Put on gown, if applicable.
 - Fully cover your torso from your neck to knees and your arms to the end of your wrists, then tie at the back.
 - The gown should be large enough to allow unrestricted movement without gaping.
 - Fasten at the back of the neck and waist.
4. Put on surgical mask or KN95 (if applicable).
 - Fit flexible band to the nose bridge.
 - Fit mask snug to face and below the chin.
5. Put on protective eyewear or face shield (if applicable).
 - Place over eyes/face and adjust to fit.
6. Put on gloves.
 - Extend the gloves to cover the wrist of the gown.



Donning must be performed in the correct order to prevent infection transmission.

Safe Practices to Remember

- Limit the number of surfaces touched when wearing PPE
- Do not touch your face while wearing PPE
- If you are interacting in close proximity to more than one individual for direct care, your PPE must be changed, and hand hygiene must be practiced in between each individual.
- If at any point your gloves become contaminated, you must dispose of them, perform hand hygiene, and then replace them with new gloves.
- Hair should be pulled back in settings where PPE is necessary.
- Jewelry should be minimized.

Doffing (removing) PPE

Following a correct doffing procedure is especially crucial in the control and prevention of infection. It is the most important step of preventing infection transmission. The doffing of PPE should protect your clothing, skin, and mucous membranes from contamination.

Remember that all PPE is contaminated after use. Perform hand hygiene immediately after each step of doffing. If you are in the isolation space, your gloves and gown (if applicable) should be removed *before* exiting the room.

1. Remove gloves.
 - Using one hand, grasp the palm of the other hand and peel off the first glove.
 - Hold the removed glove in the gloved hand.
 - Slide fingers of the ungloved hand under the remaining glove at the wrist and peel it off over the first glove.
 - Discard gloves in a waster container.
 - Perform hand hygiene appropriately
2. Remove gown.
 - Unfasten the ties, ensuring the sleeves do not make contact with your body.
 - Pull the gown away from the neck and shoulders, touching the inside only.

- Turn the gown inside out.
- Fold or roll the gown into a bundle and discard in the waste container.
- 3. Perform hand hygiene.
- 4. If applicable, exit isolation space.
- 5. Remove goggles/face shield, if applicable.
 - Remove from the back of the head by lifting headband or earpieces.
 - If reusable, place in the designated reprocessing receptacle. If not, discard in waste container.
- 6. Perform hand hygiene.
- 7. Remove mask.
 - Grasp the elastic ties looped around the ears and remove without touching the front of the mask.
 - Discard in the waste container.
- 8. Immediately perform hand hygiene.

(Adapted from diagrams by Queensland Department of Health and the CDC)

Video Resources

[Donning PPE \(CDC\)](#)

[Doffing PPE \(CDC\)](#)

[Handwashing \(CDC\)](#)